

The locally covariant Dirac field

Ko Sanders*

Institute of Theoretical Physics
University of Göttingen,
Friedrich-Hund-Platz 1, D-37077 Göttingen
and
Courant Research Centre
"Higher Order Structures in Mathematics",
University of Göttingen

November 9, 2009

Abstract

We describe the free Dirac field in a four dimensional spacetime as a locally covariant quantum field theory in the sense of Brunetti, Fredenhagen and Verch, using a representation independent construction. The freedom in the geometric constructions involved can be encoded in terms of the cohomology of the category of spin spacetimes. If we restrict ourselves to the observable algebra the cohomological obstructions vanish and the theory is unique. We establish some basic properties of the theory and discuss the class of Hadamard states, filling some technical gaps in the literature. Finally we show that the relative Cauchy evolution yields commutators with the stress-energy-momentum tensor, as in the scalar field case.

1 Introduction

Quantum field theory in curved spacetime is relevant for several purposes, such as the construction of cosmological models and to obtain a better understanding of quantum field theory in Minkowski spacetime. In order to achieve this goals in a more realistic setting it is important to go beyond the well-studied free scalar field. In this paper we will present a proof, already contained in [36], of the fact that the free Dirac field in a four dimensional globally hyperbolic spacetime can be described as a locally covariant quantum field theory in the sense of [4].

Our presentation of the Dirac field is representation independent and we emphasise categorical methods throughout in order to point out an interesting problem concerning the uniqueness of the theory. The obstruction for the definition of a unique theory can be formulated in terms of the cohomology of the category of spacetimes with a spin structure, in particular its first Stiefel-Whitney class. It seems difficult to compute this class for a category, but we will show that a unique theory can always be obtained by restriction to the observable algebras generated by even polynomials in the field, in which case the cohomological obstructions vanish.

Hadamard states can be defined in terms of a series expansion of their two-point distribution, detailing their local singularity structure. Alternatively, they can be characterised by a microlocal condition. The equivalence of these two definitions has been investigated by several authors using different techniques of proof, but in our opinion none of these arguments has been fully convincing. In our discussion we hope to close any remaining gaps in the different proofs and establish the equivalence on firm ground.

We also compute the relative Cauchy evolution of this field and obtain commutators with the stress-energy-momentum tensor, in complete analogy with the scalar field case ([4]). For this we

*E-mail: jacobus.sanders@theorie.physik.uni-goe.de

use a point-splitting procedure to renormalise the stress-energy-momentum tensor. Because we only need commutators with this tensor we do not need to treat the so-called trace anomaly, a divergent multiple of the identity operator, in detail. We refer the interested reader to [10], who also construct the extended algebra of Wick powers, relevant for perturbation theory.

The contents of this paper are organised as follows. In section 2 we review some of the mathematical background material that we need in order to describe the Dirac field. This includes first of all the Dirac algebra and the Spin group, followed by a categorical formulation of some of the differential geometry that we will need. In section 3 we describe the classical free Dirac field, starting with the geometric and algebraic aspects in subsections 3.1 and 3.2 and the equations of motion and their fundamental solutions in subsection 3.3. We discuss the uniqueness of the functorial constructions and their cohomological obstructions in subsection 3.4. We then proceed to the quantum Dirac field in section 4. In subsection 4.1 we quantise the classical Dirac field in a local and covariant way and collect some of its basic properties. Subsection 4.2 deals with Hadamard states and includes a discussion of the existing results concerning the equivalence of the microlocal and the series expansion definitions. For this purpose we also refer to appendix A, which contains several relevant and useful (but expected) results in microlocal analysis. Subsection 4.3 contains our discussion of the relative Cauchy evolution of the free Dirac field, obtaining commutators with the stress-energy-momentum tensor, but the proof of our main result there is deferred to appendix B, because it consists of rather involved computations. Finally we end with some conclusions.

Our presentation of locally covariant quantum field theory is based on the original [4] and on [16]. For the Dirac field in curved spacetime we largely follow [14] and [17], as well as our earlier [36]. For results on Clifford algebras we refer to [27] (see also [8] for a short review).

2 Mathematical preliminaries

To prepare for our discussion of the locally covariant Dirac field we present in the current section some mathematical preliminaries concerning the Dirac algebra, the Spin group and a categorical formulation of relevant aspects of differential geometry. These merely serve to fix our notation and set the scene for the subsequent sections. We also point out the relations with some other definitions and conventions in the literature.

2.1 The Dirac algebra and the Spin group

The Spin group can be embedded in the Clifford algebra of Minkowski spacetime, which we call the Dirac algebra. Therefore we will first briefly recall some results on Clifford algebras, for which we refer to [27] (note the difference in sign convention in the Clifford multiplication).

Let $\mathbb{R}^{r,s}$ be a finite dimensional real vector space with dimension $n = r + s$ and with a non-degenerate bilinear form g_{ab} which has r positive and s negative eigenvalues. The *Clifford algebra* $Cl_{r,s}$ is defined as the $\mathbb{R}$ -linear associative algebra generated by a unit element I and an orthonormal basis e_a of $\mathbb{R}^{r,n-r}$ subject to the relations:

$$e_a e_b + e_b e_a = 2g_{ab} I.$$

This definition is independent of the choice of basis. We may identify $\mathbb{R}^{r,s} \subset Cl_{r,s}$ as the subspace of monomials in the basis e_a of degree 1. The even, respectively odd, subspace of this Clifford algebra is the one spanned by monomials of even, respectively odd, degree in the basis vectors and is denoted by $Cl_{r,s}^0$, respectively $Cl_{r,s}^1$. Note that the even subspace is also a subalgebra. In the following we will be especially interested in Minkowski spacetime, $M_0 := \mathbb{R}^{1,3}$, where the bilinear form is $\eta = \text{diag}(1, -1, -1, -1)$ and where we choose an orthonormal basis g_a , $a = 0, 1, 2, 3$ with $\|g_0\|^2 = 1$, $\|\cdot\|^2$ denoting the Minkowski pseudo-norm squared. The associated Clifford algebra is called the *Dirac algebra* $D := Cl_{1,3}$ and it is characterised by

$$g_a g_b + g_b g_a = 2\eta_{ab} I. \tag{1}$$

As a vector space the Clifford algebra is naturally isomorphic to the exterior algebra. This motivates the term *volume form* for the element $g_5 := g_0 g_1 g_2 g_3$ (or in general $e := e_1 \cdots e_{r+s}$). Note the following properties:

Lemma 2.1 *We have $g_5^2 = -I$ and $g_5 v g_5^{-1} = -v$ for all $v \in M_0$. More generally, if $u \in M_0$ has $u^2 = \|u\|^2 I \neq 0$, then $u^{-1} = \frac{1}{\|u\|^2} u$ and $v \mapsto -uvu^{-1}$ defines a reflection of M_0 in the hyperplane perpendicular to u .*

Proof. These equalities follow directly from (1). For the last claim, e.g., we compute:

$$-uvu^{-1} = v - (uv + vu)u^{-1} = v - \frac{2\langle u, v \rangle}{\|u\|^2} u, \quad v \in M_0.$$

□

Standard arguments with Clifford algebras [27] give:

$$D = Cl_{1,3} \simeq Cl_{1,4}^0 \simeq Cl_{4,1}^0, \quad Cl_{4,1} \simeq M(4, \mathbb{C}),$$

where $M(4, \mathbb{C})$ denotes the algebra of complex 4×4 -matrices. In fact, $Cl_{4,1}$ is generated by the generators g_a of D together with a central element ω , corresponding to $iI \in M(4, \mathbb{C})$. Hence:

$$M(4, \mathbb{C}) \simeq \mathbb{C} \otimes_{\mathbb{R}} D. \quad (2)$$

This also implies that the center of D is spanned by I (over $\mathbb{R}$). The following Fundamental Theorem provides all the essential information we need on the Dirac algebra (for an elementary algebraic proof we refer to Pauli [31].):

Theorem 2.2 (Fundamental Theorem) *The Dirac algebra D is simple and has a unique irreducible complex representation (i.e. an $\mathbb{R}$ -linear representation $\pi: D \rightarrow M(n, \mathbb{C})$), up to equivalence. This is the representation $\pi_0: D \rightarrow M(4, \mathbb{C})$ determined by $\pi_0(g_a) = \gamma_a$ with the Dirac matrices*

$$\gamma_0 := \begin{pmatrix} O & I \\ I & 0 \end{pmatrix}, \quad \gamma_i := \begin{pmatrix} O & -\sigma_i \\ \sigma_i & 0 \end{pmatrix},$$

where σ_i are the Pauli matrices $\sigma_1 := \begin{pmatrix} O & 1 \\ 1 & 0 \end{pmatrix}$, $\sigma_2 := \begin{pmatrix} O & -i \\ i & 0 \end{pmatrix}$ and $\sigma_3 := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$.

The equivalence with another irreducible complex representation π of D is implemented by $\pi(S) = L\pi_0(S)L^{-1}$ for all $S \in D$, where $L \in GL(4, \mathbb{C})$ is unique up to a non-zero complex factor.

Consequently, for every set of matrices $\gamma'_a \in M(4, \mathbb{C})$ satisfying equation (1) there is an $L \in GL(4, \mathbb{C})$, unique up to a non-zero complex constant, such that $\gamma'_a = L\gamma_a L^{-1}$.

Proof. One can show [27] that $D \simeq M(2, \mathbb{H})$, which is simple because it is a full matrix algebra. The given matrices γ_a satisfy the Clifford relations (1) and therefore extend to a representation of D in $M(4, \mathbb{C})$.

Any complex representation $\pi: D \rightarrow M(n, \mathbb{C})$ extends to a complex representation $\tilde{\pi}$ of $M(4, \mathbb{C})$, using equality (2) and the trivial center of D , which is irreducible if π is irreducible. As $M(4, \mathbb{C})$ has only one irreducible representation up to equivalence (see [41]) this determines π up to equivalence, as stated. If $K, L \in GL(4, \mathbb{C})$ are two matrices which implement the same equivalence, then KL^{-1} commutes with D and hence $K = cL$, where $c \in \mathbb{C}$ is non-zero because K is invertible. Note that $\pi'(g_a) := \gamma'_a$ extends to a complex representation of D in $M(4, \mathbb{C})$ which is faithful (as D is simple). The last statement then follows from the previous one. □

For notational convenience we define $\gamma_5 := \pi_0(g_5)$.

We can define a determinant and trace function on D by $\det S = \det \pi(S)$ and $Tr(S) = Tr(\pi(S))$ for all $S \in D$, where π is any irreducible complex representation of D . This is well-defined by the Fundamental Theorem. The following lemma is often useful in computations:

Lemma 2.3 *We have $Tr(g_a g_b) = 4\eta_{ab}$ and $Tr([g_b, g_c] g_d g_a) = 8(\eta_{cd}\eta_{ba} - \eta_{bc}\eta_{da})$.*

Proof. Using the cyclicity of the trace and the relations (1) we find: $Tr(g_ag_b) = \frac{1}{2}Tr(g_ag_b + g_bg_a) = Tr(\eta_{ab}I) = 4\eta_{ab}$ and

$$\begin{aligned} Tr([g_b, g_c]g_dg_a) &= Tr(g_b[g_c, g_dg_a]) = Tr(g_b\{g_c, g_d\}g_a - g_bg_d\{g_c, g_a\}) \\ &= 2Tr(\eta_{cd}g_bg_a - g_bg_d\eta_{ca}) = 8(\eta_{cd}\eta_{ba} - \eta_{bd}\eta_{ca}). \end{aligned}$$

□

We now turn to the Spin group, which is the universal double covering group of the special Lorentz group and which can be constructed in an elegant way inside the Dirac algebra.

Definition 2.4 *The Pin and Spin groups of $Cl_{r,s}$ are defined as*

$$Pin_{r,s} := \{S \in Cl_{r,s} \mid S = u_1 \cdots u_k, \quad u_i \in \mathbb{R}^{r,s}, \quad u_i^2 = \pm I\},$$

$$Spin_{r,s} := Pin_{r,s} \cap Cl_{r,s}^0.$$

We let $Spin_{1,3}^0$ denote the connected component of $Spin_{1,3}$ which contains the identity.

We also define the Lorentz group $\mathcal{L} := O_{1,3}$, the special Lorentz group $\mathcal{L}_+ := SO_{1,3}$ and the special orthochronous Lorentz group $\mathcal{L}_+^\uparrow := SO_{1,3}^0$, which is the connected component of $\mathcal{L}_+$ containing the identity.

The special orthochronous Lorentz group preserves the orientation and time-orientation. For $S \in Pin_{1,3}$ the map $v \mapsto SvS^{-1}$ on M_0 is a product of reflections (up to a sign) by Lemma 2.1. Together with the fact that $\det u = \|u\|^4$ for all $u \in M_0$ this gives rise to another useful characterisation of the group $Pin_{1,3}$, which we shall not prove:¹

Proposition 2.5 $Pin_{1,3} = \{S \in D \mid \det S = 1, \forall v \in M_0 SvS^{-1} \in M_0\}.$

It can be seen from Proposition 2.5 that $Pin_{1,3}$ and $Spin_{1,3}$ are indeed Lie groups. For the universal double covering homomorphism Λ between $Pin_{1,3}$ and the Lorentz group we have the following formulae:²³

Proposition 2.6 *The map $\Lambda: Pin_{1,3} \rightarrow \mathcal{L}$ defined by $S \mapsto \Lambda_b^a(S) \in M(4, \mathbb{R})$ such that $Sg_bS^{-1} = g_a\Lambda_b^a(S)$ is the universal double covering homomorphism of Lie groups, which restricts to the universal double covering homomorphism $Spin_{1,3}^0 \rightarrow \mathcal{L}_+^\uparrow$. We have $\Lambda_b^a(S) = \frac{1}{4}Tr(g^aSg_bS^{-1})$ and the inverse of the derivative $d\Lambda: spin_{1,3}^0 \rightarrow l_+^\uparrow$ at $S = I$ is given by:*

$$(d\Lambda)^{-1}(\lambda^b_a) = \frac{1}{4}\lambda^b_ag_bg^a.$$

Proof. For the first sentence we refer to [27] Theorem 2.10 and subsequent remarks. Using the Clifford relations (1) we see that

$$\begin{aligned} \Lambda_b^a(S) &= \frac{1}{4}\eta^{ac}Tr(\eta_{cd}\Lambda_b^d(S)I) = \frac{1}{8}\eta^{ac}Tr((g_cg_d + g_dg_c)\Lambda_b^d(S)) \\ &= \frac{1}{4}\eta^{ac}Tr(g_cg_d\Lambda_b^d(S)) = \frac{1}{4}Tr(g^aSg_bS^{-1}). \end{aligned}$$

¹The definition of the Spin group in [7] corresponds to our group $Pin_{1,3}$. In [14] and [17] one uses the term Spin group for the group

$$\mathcal{S} := \{S \in M(4, \mathbb{C}) \mid \det S = 1, \quad SvS^{-1} \in M_0 \text{ for all } v \in M_0\}.$$

Note that this group cannot give a double covering of the Lorentz group, as claimed in [14] (but not in [17]), because for any $S \in \mathcal{S}$ the matrices $iS, -S, -iS$ are in $\mathcal{S}$ too. Its usefulness is based on its simple definition and the fact that $\mathcal{S}^0 = Spin_{1,3}^0$.

²These results are well-known, but we record them for definiteness to correct a sign error in the spin connection (5) that has occurred in [14, 17, 11].

³Lower case Latin indices are raised and lowered with η^{ab} , resp. η_{ab} throughout.

Expanding $\Lambda(S + \epsilon s + O(\epsilon^2))$ up to second order in ϵ we find $d\Lambda(s)^a_b = \frac{1}{4}Tr([g_b, g^a] s)$. We check that $L(\lambda^b_a) := \frac{1}{4}\lambda^b_a g_b g^a$ is an inverse of $d\Lambda$:

$$\begin{aligned} d\Lambda(L(\lambda^d_e))^a_b &= \frac{1}{16}\eta^{ac}\eta^{ef}\lambda^d_e Tr([g_b, g_c] g_d g_f) = \frac{1}{2}\eta^{ac}\eta^{ef}\lambda^d_e (\eta_{cd}\eta_{bf} - \eta_{bd}\eta_{cf}) \\ &= \frac{1}{2}(\lambda^a_b - \eta^{ae}\eta_{bd}\lambda^d_e) = \lambda^a_b, \end{aligned}$$

where we used Lemma 2.3 and the symmetry properties of $\lambda^d_e \in l_+^\uparrow$ in the last line. $\square$

2.2 Some category theory and differential geometry

The language of locally covariant quantum field theory uses category theory to express the physical ideas of locality and covariance. Any object or construction that is extended from a single spacetime (usually Minkowski spacetime) to the categorical framework gets the adjective "locally covariant". The essence of local covariance seems to have a geometric origin and, because the Dirac field in curved spacetimes involves a substantial amount of geometric constructions, it will be convenient to present the relevant differential geometry in a categorical setting here. We refrain from the urge to call this "locally covariant differential geometry", which appears to be a pleonasm.

A *category* $\mathfrak{C}$ consists of a set of objects c and a set of morphisms or arrows⁴ $\gamma : c_1 \rightarrow c_2$ between objects of $\mathfrak{C}$, such that the composition of morphisms, when defined, is associative and each object admits an identity morphism (we refer to [29] for more details). A (*covariant*) *functor* $\mathbf{F} : \mathfrak{C} \rightarrow \mathfrak{B}$ is a map between categories, which maps objects c to objects $\mathbf{F}(c)$ and morphisms $\gamma : c_1 \rightarrow c_2$ to morphisms $\mathbf{F}(\gamma) : \mathbf{F}(c_1) \rightarrow \mathbf{F}(c_2)$ such that an identity morphism maps to an identity morphism and the composition of morphisms is preserved. A *contravariant functor* $\mathbf{F} : \mathfrak{C} \rightarrow \mathfrak{B}$ is defined similarly, but reverses the direction of the morphisms: $\mathbf{F}(\gamma) : \mathbf{F}(c_2) \rightarrow \mathbf{F}(c_1)$. A *natural transformation* $t : \mathbf{F} \Rightarrow \mathbf{G}$ between covariant functors $\mathbf{F} : \mathfrak{C} \rightarrow \mathfrak{B}$ and $\mathbf{G} : \mathfrak{C} \rightarrow \mathfrak{B}$ is a map which assigns to each object c a morphism $t(c)$ of $\mathfrak{B}$, called the *component* of t at c , such that for every morphism $\gamma : c_1 \rightarrow c_2$ of $\mathfrak{C}$ we have $t(c_2) \circ \mathbf{F}(\gamma) = \mathbf{G}(\gamma) \circ t(c_1)$, which can be depicted as a commutative diagram. When a natural transformation t admits another natural transformation s such that $t(c) \circ s(c) = \text{id}_c = s(c) \circ t(c)$ for all objects c , then t is called a *natural equivalence*. In this case we write $t : \mathbf{F} \Leftrightarrow \mathbf{G}$. A natural transformation between contravariant functors or between a covariant and a contravariant functor is defined similarly, except that some arrows in the commutative diagram are reversed.

A subcategory $\mathfrak{B}$ of $\mathfrak{C}$ consists of a subset of the objects of $\mathfrak{C}$ and a subset of its morphisms in such a way that $\mathfrak{B}$ still satisfies the axioms of a category. In our case all categories will be concrete, i.e. the objects will be sets with a certain structure and the morphisms will be maps between sets. The identity morphism will always be the identity map and the composition of maps, when defined, is automatically associative. In short, our categories will be subcategories of the category $\mathbf{Set}$, whose objects are sets⁵ and whose morphisms are maps.

For our discussion of differential geometry we start with the following

Definition 2.7 *The category $\mathbf{Man}^n$ of smooth manifolds is the category whose objects are C^∞ manifolds $\mathcal{M}$ of (finite) dimension n and whose morphisms are C^∞ embeddings $\mu : \mathcal{M}_1 \rightarrow \mathcal{M}_2$.*

The category $\mathbf{Bund}'$ of fiber bundles is the category whose objects are smooth fiber bundles $p : \mathcal{B} \rightarrow \mathcal{M}$ over objects $\mathcal{M}$ of $\mathbf{Man}^n$ with bundle projection map p , and whose morphisms are C^∞ maps $\beta : \mathcal{B}_1 \rightarrow \mathcal{B}_2$ covering a morphism $\mu : \mathcal{M}_1 \rightarrow \mathcal{M}_2$ of $\mathbf{Man}^n$, i.e. such that $p_2 \circ \beta = \mu \circ p_1$. We denote by $\mathbf{Bund}$ the subcategory whose morphisms restrict to isomorphisms of the fibers.

The categories $\mathbf{VBund}'_{\mathbb{R}}$, respectively $\mathbf{VBund}'_{\mathbb{C}}$, of real (complex) vector bundles is the subcategory of $\mathbf{Bund}'$ whose objects $\mathcal{V}$ are real (complex) vector bundles and whose morphisms $\nu : \mathcal{V}_1 \rightarrow \mathcal{V}_2$ are real (complex) linear maps of the fibers. Again we denote by $\mathbf{VBund}_{\mathbb{R}}$ and $\mathbf{VBund}_{\mathbb{C}}$ the subcategories whose morphisms restrict to isomorphisms of the fibers.

⁴It is very often convenient to depict the morphisms in a diagram as arrows between objects.

⁵See [29] for some relevant remarks concerning the foundations of set theory and the use of small sets.

We could have taken all smooth maps between manifolds as morphisms of $\mathfrak{Man}^n$ or allowed all dimensions. However, local diffeomorphisms allow us to transport more structure, which enables us to describe more of the canonical differential geometric constructions as functors. We list the most important examples below. For fiber bundles, on the other hand, it will be useful to allow maps which are not isomorphisms on the fibers.^{6,7}

1. The functor $\mathbf{T} : \mathfrak{Man}^n \rightarrow \mathfrak{VBund}_{\mathbb{R}}$ assigns to every manifold $\mathcal{M}$ the tangent bundle $T\mathcal{M}$ and to every morphism $\mu : \mathcal{M}_1 \rightarrow \mathcal{M}_2$ the differential $d\mu : T\mathcal{M}_1 \rightarrow T\mathcal{M}_2$.
2. The functor⁸ $\mathbf{T}^* : \mathfrak{Man}^n \rightarrow \mathfrak{VBund}_{\mathbb{R}}$ assigns to every manifold $\mathcal{M}$ the cotangent bundle $T^*\mathcal{M}$ and to every morphism $\mu : \mathcal{M}_1 \rightarrow \mathcal{M}_2$ the push-forward $\mu_* : T\mathcal{M}_1 \rightarrow T\mathcal{M}_2$, which is defined as $\mu_*\omega := \omega \circ d\mu^{-1}$.
3. Finite direct sums and tensor products of $\mathbf{T}$ and $\mathbf{T}^*$ can also be described as functors, by extending $d\mu$ and μ_* in the obvious way.
4. The functor $\mathbf{\Lambda}^k : \mathfrak{Man}^n \rightarrow \mathfrak{VBund}_{\mathbb{R}}$ assigns to every manifold $\mathcal{M}$ the vector bundle $\Lambda^k\mathcal{M}$ of exterior k -forms and to every morphism μ the push-forward μ_* of such forms. Similarly the functor $\mathbf{\Lambda} : \mathfrak{Man}^n \rightarrow \mathfrak{VBund}_{\mathbb{R}}$ assigns to a manifold the exterior algebra and it maps morphisms to push-forwards.
5. The functor $|\mathbf{\Lambda}^n| : \mathfrak{Man}^n \rightarrow \mathfrak{VBund}_{\mathbb{R}}$ assigns to every spacetime $\mathcal{M}$ the one dimensional trivial vector bundle of densities $|\Lambda^n\mathcal{M}|$, where n is the dimension of $\mathcal{M}$. This is the vector bundle whose fiber at $x \in \mathcal{M}$ consists of functions $d : \Lambda_x^n\mathcal{M} \rightarrow \mathbb{R}$ such that $d(r\omega) = |r|\omega$ for all $r \in \mathbb{R}$ and $\omega \in \Lambda_x^n\mathcal{M}$ (cf. [2] appendix A.3). A morphism μ is mapped to the push-forward defined by $\mu_*d := d \circ \mu^*$, where $\mu^*\omega := \omega \circ d\mu$ is the pull-back.
6. In general, for a functor $\mathbf{V} : \mathfrak{Man}^n \rightarrow \mathfrak{VBund}_{\mathbb{R}}$ with $\mathcal{M} \mapsto V\mathcal{M}$ and $\mu \mapsto \beta$, the dual functor $\mathbf{V}^* : \mathfrak{Man}^n \rightarrow \mathfrak{VBund}_{\mathbb{R}}$ assigns to every manifold $\mathcal{M}$ the dual vector bundle $V^*\mathcal{M}$ of $V\mathcal{M}$ and to each morphism $\mu : \mathcal{M}_1 \rightarrow \mathcal{M}_2$ the push-forward morphism β_* defined by $\beta_*\omega := \omega \circ \beta^{-1}$.
7. When $\mathbf{V}_i : \mathfrak{Man}^n \rightarrow \mathfrak{VBund}'_{\mathbb{R}}$ for $i = 1, \dots, n$ map $\mathcal{M}$ to vector bundles over $\mathcal{M}$ one can construct the direct sum $\oplus_{i=1}^n \mathbf{V}_i$ and the tensor product $\otimes_{i=1}^n \mathbf{V}_i$.
8. Given a functor $\mathbf{V} : \mathfrak{Man}^n \rightarrow \mathfrak{VBund}_{\mathbb{R}}$ such that $V\mathcal{M}$ is a vector bundle over $\mathcal{M}$, the canonical pairing of $V\mathcal{M}$ and $V^*\mathcal{M}$ becomes a natural transformation $\langle, \rangle : \mathbf{V}^* \otimes \mathbf{V} \Rightarrow \mathbf{\Lambda}^0$ whose components cover the identity morphism.
9. For every functor $\mathbf{V} : \mathfrak{Man}^n \rightarrow \mathfrak{VBund}'_{\mathbb{R}}$ and every $r \in \mathbb{R}$ there is a natural transformation $m_r : \mathbf{V} \Leftrightarrow \mathbf{V}$ whose component at $\mathcal{M}$ is given by the map $m_r : V\mathcal{M} \rightarrow V\mathcal{M}$ such that $m_r(v) := rv$. If $r \neq 0$ this is a natural equivalence.
10. All of the functors above can be complexified, which yields functors into $\mathfrak{VBund}_{\mathbb{C}}$ or $\mathfrak{VBund}'_{\mathbb{C}}$. The complexification of $\mathbf{V}$ will be denoted by $\mathbf{V}_{\mathbb{C}}$ and there is a natural equivalence $\bar{\cdot} : \mathbf{V}_{\mathbb{C}} \Leftrightarrow \mathbf{V}_{\mathbb{C}}$ in $\mathfrak{VBund}_{\mathbb{R}}$ (or $\mathfrak{VBund}'_{\mathbb{R}}$) which sends each section to its complex conjugate.
11. The above constructions (dual, direct sum, tensor product) and natural transformations (pairing, m_r) can also be applied directly to complex vector bundles in a canonical (Hermitian) way.

⁶The unprimed categories, whose morphisms are isomorphisms of the fibers, can be described as fibered categories over $\mathfrak{Man}^n$, cf. [30] p.44.

⁷The functors $\mathbf{B} : \mathfrak{Man}^n \rightarrow \mathfrak{Bund}'$ below are all of a special type, namely they associate to a manifold $\mathcal{M}$ a fiber bundle whose base space is again $\mathcal{M}$. Although we will only use functors of this type when describing the Dirac field, the restriction is not technically necessary in our definitions.

⁸It is tempting to think of a *contravariant* functor that maps manifolds to their cotangent bundles and morphisms μ to the pull-back, $\mu^*\omega := \omega \circ d\mu$, which indeed reverses the directions of arrows and changes the order of compositions. However, the pull-back is only defined on the image of μ , so in general this does not define a morphism in $\mathfrak{VBund}'_{\mathbb{R}}$.

It will be convenient to consider distributions and integration in a categorical setting too:

Definition 2.8 $\mathfrak{TVec}$ is the category of topological vector spaces with injective continuous linear maps as morphisms. The functor $\mathbb{C} : \mathfrak{Man}^n \rightarrow \mathfrak{TVec}$ is the constant functor $\mathbb{C}$, i.e. it assigns to each object the one dimensional space $\mathbb{C}$ and to each morphism the identity morphism.

The functor of test-sections is the functor $\mathbf{C}_0^\infty : \mathfrak{VBund}'_{\mathbb{C}} \rightarrow \mathfrak{TVec}$ which maps each complex vector bundle $\mathcal{V}$ to the space $C_0^\infty(\mathcal{V})$ of compactly supported smooth sections of $\mathcal{V}$ in the test-section topology.⁹ A morphism ν , covering a morphism μ , is mapped to the push-forward ν_* defined by $\nu_*(f) = \nu \circ f \circ \mu^{-1}$ on $\mu(\mathcal{M}_1)$, extended by 0 to all of $\mathcal{M}_2$.

The functor of smooth sections is the contravariant functor $\mathbf{C}^\infty : \mathfrak{VBund}_{\mathbb{C}} \rightarrow \mathfrak{TVec}$ which maps each complex vector bundle $\mathcal{V}$ to the space $C^\infty(\mathcal{V})$ of smooth sections of $\mathcal{V}$ in the usual topology. A morphism ν , covering a morphism μ , is mapped to the pull-back ν^* defined by $\nu^*(f) = \nu^{-1} \circ f \circ \mu$.

The functor of distributions is the contravariant functor $\mathbf{Distr} : \mathfrak{VBund}'_{\mathbb{C}} \rightarrow \mathfrak{TVec}$ which maps each complex vector bundle $\mathcal{V}$ to the space $(C_0^\infty(\mathcal{V}))'$ of distributions on $\mathcal{V}$ with the weak topology induced by $C_0^\infty(\mathcal{V})$. A morphism ν , covering a morphism μ , is mapped to the pull-back ν^* defined by $\nu^*u := u \circ \nu_*$.

We will not need compactly supported distributions, but they can be defined as the functor dual to $\mathbf{C}^\infty$. Notice that objects which are not compactly supported, such as smooth sections or distributions, behave contravariantly, whereas compactly supported ones behave covariantly. Also note that the pull-back of a smooth section can only be defined for morphisms that restrict to isomorphisms of the fibers. The following constructions will be of importance in section 4:

12. There is a natural transformation $\int : \mathbf{C}_0^\infty \circ |\Lambda^n| \Rightarrow \mathbb{C}$ which assigns to each $\omega \in C_0^\infty(|\Lambda^n \mathcal{M}|)$ the integral $\int_{\mathcal{M}} \omega$.
13. Let $\mathbf{f} : \mathfrak{VBund}_{\mathbb{C}} \rightarrow \mathfrak{VBund}'_{\mathbb{C}}$ be the forgetful functor. For any functor $\mathbf{V} : \mathfrak{Man}^n \rightarrow \mathfrak{VBund}_{\mathbb{C}}$ there is a canonical natural transformation $\kappa : \mathbf{C}_0^\infty \circ \mathbf{f} \circ \mathbf{V} \Rightarrow \mathbf{C}^\infty \circ \mathbf{V}$, whose components are the canonical injections $C_0^\infty(V\mathcal{M}) \subset C^\infty(V\mathcal{M})$.
14. For any functor $\mathbf{V} : \mathfrak{Man}^n \rightarrow \mathfrak{VBund}_{\mathbb{C}}$ there is a canonical natural transformation $\iota : \mathbf{C}^\infty \circ (\mathbf{V} \otimes |\Lambda^n|) \Rightarrow \mathbf{Distr} \circ \mathbf{f} \circ \mathbf{V}^*$ given by $\iota_{\mathcal{M}}(f \otimes \omega) := \int_{\mathcal{M}} \langle \cdot, f \rangle \omega$ for any smooth section f of $V\mathcal{M}$ and any density ω on $\mathcal{M}$. Each component of ι is injective.
15. Given a pair of functors $\mathbf{V}_i : \mathfrak{Man}^n \rightarrow \mathfrak{VBund}'_{\mathbb{C}}$, $i = 1, 2$, any natural transformation $t : \mathbf{V}_1 \Rightarrow \mathbf{V}_2$ lifts to a corresponding natural transformation $T : \mathbf{C}_0^\infty \circ \mathbf{V}_1 \Rightarrow \mathbf{C}_0^\infty \circ \mathbf{V}_2$ defined pointwise by $T_{\mathcal{M}}f := t_{\mathcal{M}} \circ f$. For $\mathbf{V}_i : \mathfrak{Man}^n \rightarrow \mathfrak{VBund}_{\mathbb{C}}$ and a natural transformation $t : \mathbf{V}_1 \Rightarrow \mathbf{V}_2$ we can similarly define $T : \mathbf{C}^\infty \circ \mathbf{V}_1 \Rightarrow \mathbf{C}^\infty \circ \mathbf{V}_2$ pointwise by $T_{\mathcal{M}}f := t_{\mathcal{M}} \circ f$.

Where convenient we will often identify a functor $\mathbf{V} : \mathfrak{Man}^n \rightarrow \mathfrak{VBund}_{\mathbb{C}}$ with the functor $\mathbf{f} \circ \mathbf{V}$, omitting the forgetful functor, as this rarely leads to confusion.

Next we add the structure of a semi-Riemannian metric:

Definition 2.9 The category $\mathfrak{SRMan}^n$ of semi-Riemannian manifolds is the subcategory of $\mathfrak{Man}^n$ whose objects $M = (\mathcal{M}, g)$ are C^∞ manifolds $\mathcal{M}$ of dimension n with a semi-Riemannian metric g and whose morphisms $m : M_1 \rightarrow M_2$ are given by the isometric morphisms in $\mathfrak{Man}^n$, i.e. morphisms $\mu : \mathcal{M}_1 \rightarrow \mathcal{M}_2$ such that $\mu^*g_1 = g_2|_{\mu(\mathcal{M}_1)}$.

The extra structure gives rise to extra functors and natural equivalences that are of interest to us:

16. The forgetful functor $\mathbf{f} : \mathfrak{SRMan}^n \rightarrow \mathfrak{Man}^n$ assigns to each $M = (\mathcal{M}, g)$ the underlying manifold $\mathcal{M}$ and to each morphism m the underlying morphism μ in $\mathfrak{Man}^n$.
17. We will write $\mathbf{T}$, respectively $\mathbf{T}^*$, for the functors $\mathbf{T} \circ \mathbf{f}$, respectively $\mathbf{T}^* \circ \mathbf{f}$, from $\mathfrak{SRMan}^n$ to $\mathfrak{VBund}_{\mathbb{R}}$. There is then a natural equivalence $G : \mathbf{T} \Leftrightarrow \mathbf{T}^*$ whose component at $M = (\mathcal{M}, g)$ is given by the map $G_{\mathcal{M}} : T\mathcal{M} \rightarrow T^*\mathcal{M}$ such that $v \mapsto g(v, \cdot)$.

⁹For a precise definition of the well-known topologies on test-sections and smooth sections we refer to [13] Ch. 17.

18. The functor $\mathbf{F} : \mathfrak{SMan}^n \rightarrow \mathfrak{VBund}_{\mathbb{R}}$ assigns to each object M the frame bundle FM , i.e. the bundle whose fiber at a point $x \in \mathcal{M}$ consists of all orthonormal bases of $T_x\mathcal{M}$ in the metric g . This fiber is a subset of $T^{\otimes n}\mathcal{M}$. A morphism m is mapped to the push-forward μ_* acting on $FM \subset T^{\otimes n}\mathcal{M}$.
19. The functor $\mathbf{Cl} : \mathfrak{SMan}^n \rightarrow \mathfrak{VBund}_{\mathbb{R}}$ assigns to each object $M = (\mathcal{M}, g)$ the Clifford bundle $Cl\mathcal{M}$, which is the vector bundle whose fiber at $x \in \mathcal{M}$ is the Clifford algebra of $(T_x\mathcal{M}, g)$ viewed as a linear space. A morphism m is mapped to the push-forward acting on $Cl\mathcal{M} \subset \oplus_{k=0}^n T^{\otimes k}\mathcal{M}$. Note that $\mathbf{Cl}$ is naturally equivalent to $\mathbf{\Lambda} \circ \mathbf{f}$, because we ignore the algebraic structure on these vector bundles.
20. We define the *volume form functor* $\mathbf{vol} : \mathfrak{SMan}^n \rightarrow \mathfrak{VBund}_{\mathbb{R}}$ as $\mathbf{vol} := |\mathbf{\Lambda}^n| \circ \mathbf{f}$. When $m : M_1 \rightarrow M_2$ is a morphism and $d\text{vol}_i := \sqrt{|\det g_i|}$ the metric induced volume form on M_i , then $\mathbf{vol}$ maps $d\text{vol}_1$ to the restriction of $d\text{vol}_2$ to $m(\mathcal{M}_1)$. There is a canonical natural equivalence from $\mathbf{\Lambda}^0$ to $\mathbf{vol}$, which consists of multiplication with the metric induced volume form.
21. Similarly there are natural equivalences between any functor $\mathbf{V} : \mathfrak{SMan}^n \rightarrow \mathfrak{VBund}_{\mathbb{C}}$ and $\mathbf{V} \otimes |\mathbf{\Lambda}^n|$. Therefore we obtain a canonical natural transformation $\iota : \mathbf{C}^\infty \circ \mathbf{V} \Rightarrow \mathbf{Distr} \circ \mathbf{V}^*$, the components of which are injective.

3 The classical Dirac field

After these mathematical preliminaries we are now ready to start constructing the classical free Dirac field (as a locally covariant classical field). We will first describe the geometric and algebraic constructions, before we discuss the Dirac equation and its fundamental solutions. We close by investigating to what extent the relations between the Dirac operator, charge conjugation and adjoint map fix the structure of the theory and find that the non-uniqueness can be characterised in terms of the cohomology of the category of spin spacetimes.

3.1 Geometric aspects

In order to describe the Dirac field we need to introduce the notion of a spin structure on a spacetime, combining the geometric and the algebraic results of section 2. This is the purpose of the current subsection.

The systems that we will consider are intended to model Dirac quantum fields living in a (region of) spacetime which is endowed with a fixed Lorentzian metric (a background gravitational field). Mathematically these regions are modelled as follows:

Definition 3.1 *By the term globally hyperbolic spacetime we will mean a connected, Hausdorff, paracompact, C^∞ Lorentzian manifold $M = (\mathcal{M}, g)$ of dimension $d = 4$, which is oriented, time-oriented and admits a Cauchy surface.*

A subset $O \subset \mathcal{M}$ of a globally hyperbolic spacetime M is called causally convex iff for all $x, y \in O$ all causal curves in $\mathcal{M}$ from x to y lie entirely in O .

The category $\mathbf{Spac}$ is the subcategory of $\mathfrak{SMan}^n$ whose objects are all globally hyperbolic spacetimes $M = (\mathcal{M}, g)$ and whose morphisms are isometric embeddings ψ that preserve the orientation and time-orientation and such that $\psi(\mathcal{M}_1)$ is causally convex.

Most notations we use concerning the causal structure of spacetimes are standard, cf. [42]. The importance of causally convex sets is that for any morphism ψ the causal structure of M_1 coincides with that of $\psi(M_1)$ inside M_2 :

$$\psi(J_{M_1}^\pm(x)) = J_{M_2}^\pm(\psi(x)) \cap \psi(\mathcal{M}_1), \quad x \in \mathcal{M}_1.$$

If $O \subset \mathcal{M}$ is a connected open causally convex set, then $(O, g|_O)$ defines a globally hyperbolic spacetime in its own right. In this case there is a canonical morphism $I_{M,O} : O \rightarrow M$ given by

the canonical embedding $\iota: O \rightarrow \mathcal{M}$. We will often drop $I_{M,O}$ and ι from the notation and simply write $O \subset M$.

Notice that there is a forgetful functor $\mathbf{f}: \mathfrak{Spac} \rightarrow \mathfrak{SMan}^n$ and that we can define the functor $\mathbf{F}_+^\uparrow: \mathfrak{Spac} \rightarrow \mathfrak{Bund}$ of oriented, time-oriented orthonormal frames $F_+^\uparrow \mathcal{M}$ for the tangent bundle, in analogy to section 2.2. This is a principal $\mathcal{L}_+^\uparrow$ -bundle over M , where the special orthochronous Lorentz group $\mathcal{L}_+^\uparrow$ acts from the right, i.e., given $e = (x, e_0, \dots, e_3) \in F_+^\uparrow M$, where $x \in \mathcal{M}$ and $e_a \in T_x M$ such that $g_x(e_a, e_b) = \eta_{ab}$ and e_0 is future pointing, the action of Λ is defined by $R_\Lambda e = e' = (x, e'_0, \dots, e'_3)$ where $e'_a = e_b \Lambda_a^b$.

Definition 3.2 A spin structure on M is a pair (SM, π) , where SM is a principal $Spin_{1,3}^0$ -bundle over M , the spin frame bundle, with a right action R_S , $S \in Spin_{1,3}^0$, and $\pi: SM \rightarrow FM$, the spin frame projection, is a base-point preserving bundle homomorphism such that

$$\pi \circ R_S = R_{\Lambda(S)} \circ \pi,$$

where $S \mapsto \Lambda(S)$ is the universal covering map (cf. Proposition 2.6).

A globally hyperbolic spin spacetime $SM = (\mathcal{M}, g, SM, \pi)$ is an object $M = (\mathcal{M}, g)$ of $\mathfrak{Spac}$ which is endowed with the spin structure (SM, π) .

The category $\mathfrak{SSpac}$ is the subcategory of $\mathfrak{Bund}$ whose objects are all globally hyperbolic spin spacetimes $SM = (\mathcal{M}, g, SM, \pi)$ and whose morphisms $\chi: SM_1 \rightarrow SM_2$ cover a morphism $\psi: M_1 \rightarrow M_2$ in $\mathfrak{Spac}$ and satisfy $\chi \circ (R_1)_S = (R_2)_S \circ \chi$ and $\pi_2 \circ \chi = \psi_* \circ \pi_1$, where p_i are the bundle projections, π_i the spin frame projections and ψ_* the push-forward.

Note that a morphism acts as a diffeomorphism of the fibers, because it intertwines the group action.

Every globally hyperbolic spacetime admits a spin structure, which need not be unique [19, 20, 14, 27]. We will regard distinct spin structures on the same underlying spacetime as distinct spin spacetimes.¹⁰ Spinor and cospinor fields are sections of vector bundles associated to the spin frame bundle. We will require that the assignment of these vector bundles is functorial:

Definition 3.3 A locally covariant spinor bundle is a functor $\mathbf{V}: \mathfrak{SSpac} \rightarrow \mathfrak{VBund}_{\mathbb{C}}$, written as $SM \mapsto V_{SM}$, $\chi \mapsto \nu$, such that χ and ν cover the same morphism ψ in $\mathfrak{Spac}$ and such that each V_{SM} is a vector bundle associated to the spin frame bundle SM through some representation. The dual functor $\mathbf{V}^*$ is called a locally covariant cospinor bundle. Smooth sections of V_{SM} , respectively V_{SM}^* , are called (Dirac) spinors (or spinor fields), respectively cospinors (cospinor fields).

The condition in the definition of a locally covariant spinor bundle ensures that the vector bundle V_{SM} and the spin frame bundle SM are both bundles over the same spacetime M .

For definiteness we pick out the following standard choice of locally covariant spinor and cospinor bundles:

Definition 3.4 The standard locally covariant Dirac spinor bundle $\mathbf{D}_0: \mathfrak{SSpac} \rightarrow \mathfrak{VBund}_{\mathbb{C}}$ is the locally covariant spinor bundle which associates to each object SM of $\mathfrak{SSpac}$ the associated vector bundle $D_0 M = SM \times_{Spin_{1,3}^0} \mathbb{C}^4$ of SM with the representation π_0 , and which maps each morphism $\chi: SM_1 \rightarrow SM_2$ to the morphism $\xi: D_0 M_1 \rightarrow D_0 M_2$ given by $\xi([E, z]) := [\chi(E), z]$.

The standard locally covariant Dirac cospinor bundle $\mathbf{D}_0^*$ is the dual functor of $\mathbf{D}_0$.

Recall that a point in $D_0 M$ consists of an equivalence class of pairs $(E, z) \in SM \times \mathbb{C}^4$, where the equivalence is given by

$$[R_S E, z] = [E, \pi_0(S)z].$$

¹⁰There exists another approach to spinors, which considers on each spacetime the Clifford bundle. This Clifford bundle is functorial in its dependence on the spacetime, but it does not generally define a spin structure. Indeed, at each point one can identify the Spin group inside the fiber of the Clifford bundle, but there may not be any projection from these Spin groups onto the frame bundle that intertwines the actions of the structure groups, the obstruction being a topological twist. (Conversely, every spin structure can be seen as a topologically twisted copy of the Spin groups in the Clifford bundle.) Nevertheless, it appears to provide sufficient structure to describe all the relevant physics in a functorial way. We refer to [40] for more information on this approach.

The dual functor $\mathbf{D}_0^*$ then assigns to each SM the dual vector bundle D_0^*M whose points are equivalence classes of pairs $(E, w^*) \in SM \times (\mathbb{C}^4)^*$, where the equivalence is given by $[R_S E, w^*] = [E, w^* \pi_0(S^{-1})]$. (Here we consider $w^* \in (\mathbb{C}^4)^*$ as a row vector, whereas $z \in \mathbb{C}^4$ is treated as a column vector.)

For any object SM the unique connection ∇_{SM} on TM which is compatible with the metric, $\nabla_{SM}g = 0$, can be described by an $l_+^\dagger$ -valued one-form $(\Omega_{SM})_a^b$ on the orthonormal frame bundle $F_+^\dagger M$ (cf. [25] Ch.2 Proposition 1.1), where $l_+^\dagger$ is the Lie-algebra of $\mathcal{L}_+^\dagger$, which can be identified with the tangent space of the fiber of $F_+^\dagger M$ at any point. For every local section e of $F_+^\dagger M$ the pull-back $\omega_a^b := e^*(\Omega_a^b)$ consists exactly of the connection one-forms of ∇_{SM} expressed in the orthonormal frame e_a . The one-form $(\Omega_{SM})_a^b$ can be pulled back by the spin frame projection π and lifted to a $spin_{1,3}^0$ -valued one-form Σ_{SM} on SM :

$$\Sigma_{SM} := (d\Lambda)^{-1} \pi^*((\Omega_{SM})_a^b) = \frac{1}{4} p^*((\Omega_{SM})_a^b) g_b g^a,$$

where the last equality uses Proposition 2.6. The one-form Σ_{SM} determines a connection on the spin frame bundle SM . For any associated vector bundle DM we then find a connection, also denoted by ∇_{SM} , determined by the connection one-forms $\sigma := E^*(\Sigma_{SM})$ in a local section E of SM , as represented on DM (we will give an explicit expression for σ in equation (5) below). The connection can be viewed as a map $\nabla_{SM}: C_0^\infty(D_0M) \rightarrow C_0^\infty(T^*M \otimes D_0M)$, which is a component of a natural transformation¹¹ $\nabla: C_0^\infty \circ \mathbf{D}_0 \Rightarrow C_0^\infty \circ (\mathbf{T}^* \otimes \mathbf{D}_0)$. The Leibniz rule allows us to extend it to mixed spinor-tensors, using e.g. $\nabla_a \langle v, u \rangle = \langle \nabla_a v, u \rangle + \langle v, \nabla_a u \rangle$.

3.2 Adjoints, charge conjugation and the Dirac operator,

We now define the adjoint and charge conjugation maps on spinors and cospinors. These are special cases of the Fundamental Theorem 2.2, using the complex conjugate and adjoint matrices¹² (cf. [21]).

Theorem 3.5 *For any irreducible complex representation π of the Dirac algebra D there are matrices $A, C \in GL(4, \mathbb{C})$ such that*

$$\begin{aligned} A &= A^*, \quad \pi(g_a)^* = A \pi(g_a) A^{-1}, \quad A n > 0, \\ \overline{C} C &= I, \quad -\overline{\pi(g_a)} = C \pi(g_a) C^{-1} \end{aligned} \tag{3}$$

for all future pointing time-like vectors $n \in M_0 \subset D$. We have for all $S \in Spin_{1,3}^0$:

$$\begin{aligned} A &= -C^* A^T C, \\ \pi(S)^* A \pi(S) &= A, \quad \pi(S^{-1}) C^{-1} \overline{\pi(S)} = C^{-1}. \end{aligned}$$

Moreover, if $A', C' \in M(4, \mathbb{C})$ have the properties stated above for the irreducible complex representation π' of D , then there is an $L \in GL(4, \mathbb{C})$, unique up to a sign, such that $L^* A' L = A$, $(\overline{L})^{-1} C' L = C$ and $\pi = L^{-1} \pi' L$ on D .

Proof. To prove the existence of A and C in the representation π_0 we may take $A = A_0 := \gamma_0$, $C = C_0 := \gamma_2$ and check the required properties straightforwardly. Note for example that

$$\gamma_0 n^a \gamma_a = \begin{pmatrix} n^0 I + n^i \sigma_i & 0 \\ 0 & n^0 I - n^i \sigma_i \end{pmatrix} > 0,$$

¹¹Alternatively we could have written the connection as a natural transformation from the 1-jet bundle extension of $\mathbf{D}_0$ to $\mathbf{T}^* \otimes \mathbf{D}_0$.

¹²On a general representation space of complex dimension four one can define many complex conjugations and Hermitean inner products. In order to obtain the desired equalities involving adjoint and charge conjugate spinors later on we need these two operations to be compatible, i.e. $\langle \overline{v}, \overline{w} \rangle = \langle v, w \rangle$. Without loss of generality we can then use the standard complex conjugation and Hermitean inner product on $\mathbb{C}^4$.

because $\det(n^0 I \pm n^i \sigma_i) = n^2 > 0$ and $\text{Tr}(n^0 I \pm n^i \sigma_i) = 2n^0 > 0$. To prove the existence of A and C in a general irreducible complex representation π one writes $\gamma_a = K\pi(g_a)K^{-1}$ by Theorem 2.2 and verifies that $A = K^*A_0K$ and $C = \overline{K}^{-1}C_0K$ will do.

Given A', C' satisfying equation (3) for π' we can fix $K \in GL(4, \mathbb{C})$ such that $\pi' = K\pi K^{-1}$ on D and the desired matrix L must be $L = zK$ for some $z \neq 0$ by the Fundamental Theorem 2.2. Now set $\tilde{A} := K^*A'K$ and $\tilde{C} := (\overline{K})^{-1}C'K$ and note that $\tilde{A}$ and $\tilde{C}$ satisfy (3) for π . Because the sets of matrices $\pi(g_a)^*$ and $-\pi(g_a)$ both satisfy the relations (1) we must have $aA = \tilde{A}$ and $cC = \tilde{C}$ for some non-zero complex factors a and c , again by the Fundamental Theorem. Also, $|c| = 1$ because $\overline{C}C = I$ and $a > 0$ because $A = A^*$ and $A\pi(n) > 0$ for future pointing time-like vectors. Hence, $|z|^2 = a$ and $z = c\bar{z}$, which fixes z (and L) up to a sign. This proves the last statement.

The equation $A = -C^*A^TC$ holds for A_0, C_0 and therefore also in general. For a unit vector $u = u^a g_a$ we have $u^2 = \pm I$ and hence

$$\pi(u)^* A \pi(u) = u^a u^b \pi(g_a)^* A \pi(g_b) = u^a u^b A \pi(g_a g_b) = A \pi(u^2) = \pm A.$$

For $S \in Spin_{1,3}$ we must therefore have that $\pi(S)^* A \pi(S) = \pm A$, by definition of the Spin group. For $S = I$ the sign is a plus, so by continuity and connectedness we conclude that $\pi(S)^* A \pi(S) = A$ for all $S \in Spin_{1,3}^0$. For C we use the fact that

$$\pi(u^{-1})C^{-1}\overline{\pi(u)} = -\pi(u)^{-1}\pi(u)C^{-1} = -C^{-1}$$

and hence $\pi(S^{-1})C^{-1}\overline{\pi(S)} = C^{-1}$ for all $S \in Spin_{1,3}$. $\square$

Note that $g_5 \in Spin_{1,3} \setminus Spin_{1,3}^0$. Indeed, using π_0 and $A_0 = \gamma_0$ in Theorem 3.5 we see that $\gamma_5^* A_0 \gamma_5 = -A_0$, so $g_5 \in Spin_{1,3}$ by definition, but not in $Spin_{1,3}^0$.

In the following theorem we use the fact that for any pair of natural transformations $t, t' : \mathfrak{S}\mathfrak{P}\mathfrak{a}\mathfrak{c} \Rightarrow \mathfrak{V}\mathfrak{B}\mathfrak{u}\mathfrak{n}\mathfrak{d}'_{\mathbb{C}}$ we can define the sum $t + t'$ and the tensor product $t \otimes t'$ componentwise.

Theorem 3.6 *The standard locally covariant Dirac spinor and cospinor bundles admit natural ($\mathbb{C}$ -antilinear) equivalences $^+ : \mathbf{D}_0 \Leftrightarrow \mathbf{D}_0^*$, $^c : \mathbf{D}_0 \Leftrightarrow \mathbf{D}_0$, $^c : \mathbf{D}_0^* \Leftrightarrow \mathbf{D}_0^*$ in $\mathfrak{V}\mathfrak{B}\mathfrak{u}\mathfrak{n}\mathfrak{d}_{\mathbb{R}}$ and a natural transformation $\gamma : \mathbf{D}_0 \Rightarrow \mathbf{T}^* \otimes \mathbf{D}_0$ in $\mathfrak{V}\mathfrak{B}\mathfrak{u}\mathfrak{n}\mathfrak{d}'_{\mathbb{C}}$ such that all components cover the identity morphism and the following equations hold both on spinors and cospinors (i.e. we denote the inverses of $^+$ and c by the same symbol):*

$$\begin{aligned} ^+ \circ ^+ &= 1 = ^c \circ ^c & ^+ \circ ^c &= -1 \circ ^c \circ ^+ \\ \langle , \rangle \circ S \circ (^+ \otimes ^+) &= \quad ^- \circ \langle , \rangle & \langle , \rangle \circ (^c \otimes ^c) &= \quad ^- \circ \langle , \rangle \\ (1 \otimes ^+) \circ \gamma &= \gamma^* \circ ^+ & (1 \otimes ^c) \circ \gamma &= -1 \circ \gamma \circ ^c \end{aligned} \tag{4}$$

$$(1 + S \otimes 1) \circ (1 \otimes \gamma) \circ \gamma = (2 \circ g) \otimes 1,$$

$$\nabla \circ \gamma = \gamma \circ \nabla,$$

where $S : \mathbf{D}_0 \otimes \mathbf{D}_0^* \Leftrightarrow \mathbf{D}_0^* \otimes \mathbf{D}_0$ and $S : \mathbf{T}^* \otimes \mathbf{T}^* \Leftrightarrow \mathbf{T}^* \otimes \mathbf{T}^*$ swap the factors in the tensor product, $g : \Lambda^0 \Rightarrow \mathbf{T}^* \otimes \mathbf{T}^*$ maps the function 1 to the metric g and $\gamma^* : \mathbf{D}_0^* \Rightarrow \mathbf{T}^* \otimes \mathbf{D}_0^*$ is the adjoint map of γ under the canonical pairing $\langle , \rangle$. Furthermore, for every object SM , every time-like future pointing tangent vector $n \in TM$ and every $v \in D_0 M$ we have $\langle n \otimes v^+, \gamma(v) \rangle \geq 0$.

The natural transformation γ can also be seen as a natural transformation $\mathbf{T} \Rightarrow \text{End}(\mathbf{D}_0)$ or $\mathbf{T} \Rightarrow \text{End}(\mathbf{D}_0^*)$. Equations (4) simply give the usual computational rules for spinors and cospinors in a functorial setting. Thus, for every SM and every $p \in D_0 M$, $q \in D_0^* M$ we have:

$$\begin{aligned} p^{++} &= p = p^{cc} & p^{c+} &= -p^{+c} \\ \langle p^+, q^+ \rangle &= \quad \overline{\langle q, p \rangle} & \langle q^c, p^c \rangle &= \quad \langle q, p \rangle \\ (\gamma_\mu p)^+ &= p^+ \gamma_\mu & (\gamma_\mu p)^c &= -\gamma_\mu p^c \\ \gamma_\mu \gamma_\nu + \gamma_\nu \gamma_\mu &= 2g_{\mu\nu} I, & \nabla_a \gamma_b &\equiv 0, \end{aligned}$$

where we have dropped the subscript SM to lighten the notation.

Proof. The canonical pairing $\langle, \rangle : \mathbf{D}_0^* \otimes \mathbf{D}_0 \Rightarrow \mathbf{\Lambda}_{\mathbb{C}}^0$ on SM is given by $\langle [E, w^*], [E, z] \rangle = \langle w, z \rangle$, where the right-hand side is the standard Hermitean inner product on $\mathbb{C}^4$. Note that this is well-defined, because we can always get the same $E \in SM$ on the left-hand side by a suitable action of $Spin_{1,3}^0$. The components of the natural equivalences $^+$ and c on each SM are defined using the matrices A_0 and C_0 of Theorem 3.5 and their properties:

$$\begin{aligned} [E, z]^c &:= [E, C_0^{-1} \bar{z}] & [E, w^*]^c &:= [E, \overline{w^*} C_0] \\ [E, z]^+ &:= [E, z^* A_0] & [E, w^*]^+ &:= [E, A_0^{-1} w]. \end{aligned}$$

These are well-defined isomorphisms in $\mathfrak{VBund}_{\mathbb{R}}$ and they give rise to natural equivalences satisfying the first two lines of equation (4).

Now fix $E \in SM$, let e_a be the orthonormal basis $(e_0, \dots, e_3) = \pi(E)$ of $T_{p(E)}M$, where $\pi : SM \rightarrow FM$ is the spin frame projection, and let e^a be the dual basis of $T_{p(E)}^*M$. On SM we define the component of the natural transformation γ on SM to be

$$\gamma([E, z]) := e^a \otimes [E, \gamma_a z].$$

This is well-defined, because a different section $E' := R_S E$ gives rise to the frame $e'_a = e_b \Lambda^b_a(S)$ and the dual frame $(e')^a = \Lambda^a_b(S^{-1}) e^b$ and on the other hand $\pi_0(S^{-1}) \gamma_a \pi_0(S) = \gamma_b \Lambda^b_a(S^{-1})$ by definition of Λ (Proposition 2.6). γ is indeed a morphism in $\mathfrak{VBund}'_{\mathbb{C}}$ and gives rise to a natural transformation. The third line of equation (4) follows again from the properties of A and C (see Theorem 3.5):

$$\begin{aligned} \gamma([E, z]^c) &= e^a \otimes [E, \gamma_a C_0^{-1} \bar{z}] = -e^a \otimes [E, C_0^{-1} \overline{\gamma_a z}] = -(\gamma([E, z]))^c, \\ \gamma^*([E, z]^+) &= e^a \otimes [E, z^* A_0 \gamma_a] = e^a \otimes [E, z^* \gamma_a^* A] = (\gamma([E, z]))^+ \end{aligned}$$

and similarly on cospinors. Also,

$$\begin{aligned} \nabla_b \gamma_a &= \sigma_b \gamma_a - \gamma_a \sigma_b - \Gamma_{ba}^c \gamma_c = \frac{1}{4} \Gamma_{bd}^c (\gamma_c \gamma^d \gamma_a - \gamma_a \gamma_c \gamma^d) - \Gamma_{ba}^c \gamma_c \\ &= \frac{1}{4} \Gamma_{bd}^c (\gamma_c \{ \gamma^d, \gamma_a \} - \{ \gamma_a, \gamma_c \} \gamma^d - 4 \delta_a^d \gamma_c) = \frac{-1}{2} \Gamma_{bd}^c (\delta_a^d \gamma_c + \eta_{ac} \gamma^d) = 0. \end{aligned}$$

Finally, for every object SM , every future pointing tangent vector $n \in TM$ and every $v \in D_0 M$ we have $\langle n \otimes v^+, \gamma(v) \rangle = \langle v^+, A n^a \gamma_a v \rangle \geq 0$ again by Theorem 3.5. $\square$

In terms of the Christoffel symbols $\Gamma_{\mu\nu}^\rho$, the frame e_ρ^a and representing g_a on $D_0 M$ using the $\text{End}(D_0 M)$ -valued one-forms γ , the connection one-forms of the spin connection can be expressed as¹³

$$\begin{aligned} \sigma_b &:= \frac{1}{4} \Gamma_{bc}^a \gamma_a \gamma^c, \\ \Gamma_{bc}^a &= -e_c^\rho (e_b^\sigma \partial_\sigma e_\rho^a) + e_\rho^a e_b^\mu e_c^\nu \Gamma_{\mu\nu}^\rho. \end{aligned} \tag{5}$$

The Dirac operator is defined on spinors and cospinors by

$$\nabla_{SM} := \gamma^a \nabla_a.$$

This defines natural transformations $\nabla : \mathbf{C}_0^\infty \circ \mathbf{D}_0 \Rightarrow \mathbf{C}_0^\infty \circ \mathbf{D}_0$, respectively $\nabla : \mathbf{C}_0^\infty \circ \mathbf{D}_0^* \Rightarrow \mathbf{C}_0^\infty \circ \mathbf{D}_0^*$. The intertwining relations of the adjoint and charge conjugation with the Dirac operator follow from their intertwining with γ in Theorem 3.6:

Proposition 3.7 $\nabla \circ ^+ = ^+ \circ \nabla$, $\nabla \circ ^c = -1 \circ ^c \circ \nabla$.

¹³Note the sign error in [14, 17].

Proof. Recall that $^+$ and c can be defined pointwise on test-sections. Hence, on any object SM

$$\begin{aligned} (\nabla v)^c &= ((\partial_a v - v \sigma_a) \gamma^a)^c = (\partial_a \bar{v} - \bar{v} \sigma_a) \bar{\gamma}^a C \\ &= -(\partial(\bar{v} C) - \bar{v} C \sigma_a) \gamma^a = -\nabla(\bar{v} C) = -\nabla v^c, \\ (\nabla u)^+ &= (\gamma^a (\partial_a u + \sigma_a u))^+ = (\partial_a u^* + u^* \sigma_a^*) (\gamma^a)^* A \\ &= (\partial_a (u^* A) - u^* A \sigma_a) \gamma^a = \nabla(u^* A) = \nabla u^+, \end{aligned}$$

where the minus sign in the last line appears because the order of the two factors of γ in the expression for σ_a needs to be changed. It follows that $(\nabla v)^+ = (\nabla v^{++})^+ = (\nabla v^+)^{++} = \nabla v^+$ and $(\nabla u)^c = (\nabla u^+)^{+c} = -(\nabla u^+)^{c+} = (\nabla u^{+c})^+ = -(\nabla u^{c+})^+ = -\nabla u^c$. $\square$

Remark 3.8 *A change in the sign convention, $\tilde{\eta} := -\eta$, has no physical consequences. In fact, this simply gives rise to $D \simeq Cl_{3,1}$ as the Dirac algebra, but since $Cl_{3,1}^0 = Cl_{1,3}^0$ nothing changes in the representation¹⁴ of the group $Spin_{1,3}^0 = Spin_{3,1}^0$. To accommodate this change one can set $\tilde{\gamma}_a := i\gamma_a$ in equation (1), which yields the same Dirac algebra and other constructions (although we do get signs for all covectors when raising or lowering indices with $\tilde{\eta}$). This also implies that one should drop the factor i in front of the Dirac operator in the Dirac equation (6) below, which ensures that $P_c P = P P_c$ will still be a wave operator. We can also keep the same matrices A, C , which now must satisfy the relations:*

$$-\tilde{\gamma}_a^* = A \tilde{\gamma}_a A^{-1}, \quad \bar{\tilde{\gamma}}_a = C \tilde{\gamma}_a C^{-1}.$$

The spinor and cospinor bundle and the adjoint and charge conjugation maps then remain the same and all the relations between these operations and the Dirac operator remain valid.

3.3 The Dirac equation and its fundamental solutions

The Dirac equation on spinor and cospinor fields, respectively, on a spin spacetime SM is

$$(-i\nabla + m)u = 0 \quad (i\nabla + m)v = 0, \quad (6)$$

where the constant $m \geq 0$ is to be interpreted as the mass of the field. These equations can be derived as the Euler-Lagrange equations from the action $\mathcal{S}_D := \int \mathcal{L}_D$ with the Lagrangian density¹⁵

$$\mathcal{L}_D := \langle u^+, (-i\nabla + m)u \rangle d\text{vol}_g \quad (7)$$

by varying with respect to u and u^+ , viewed as independent fields. The canonical momentum of the field u on a Cauchy surface C with future pointing normal vector field n is defined as

$$\pi(x) := \frac{1}{\sqrt{-\det g(x)}} \frac{\delta \mathcal{S}_D}{\delta(n^\mu \nabla_\mu \psi(x))} = -i\psi^+(x) \not{n}(x). \quad (8)$$

We will write $P := -i\nabla + m$ for the operator on spinors and $P_c := i\nabla + m$ for the operator on cospinors. These are components of natural transformations $P: \mathbf{C}_0^\infty \circ \mathbf{D}_0 \Rightarrow \mathbf{C}_0^\infty \circ \mathbf{D}_0$, $P: \mathbf{C}^\infty \circ \mathbf{D}_0 \Rightarrow$

¹⁴Notice that a complex irreducible representation of $Cl_{1,3}$ extends to an irreducible representation of $M(4, \mathbb{C})$ and therefore also gives a complex irreducible representation of $Cl_{3,1}$ and vice versa. The standard Clifford algebra isomorphism $Cl_{3,1} \simeq M(4, \mathbb{R})$ appears if and only if the representation of $Cl_{1,3}$ is a Majorana representation, i.e. if $\bar{\gamma}_a = -\gamma_a$. In that case we also find (see e.g. [7] p.332)

$$Pin_{3,1} \simeq \{S \in M(4, \mathbb{R}) \mid \det S = 1, \forall v \in M_0 S v S^{-1} \in M_0\} \neq Pin_{1,3}.$$

¹⁵The Lagrangian is a natural transformation between the functor $\mathbf{J}_1 \mathbf{D}_0$, which assigns to each spin spacetime SM the first-order jet bundle $J_1 D_0 M$ of the spinor bundle $D_0 M$, to the functor $|\mathbf{\Lambda}^n|$ of densities. A component of this natural transformation covers the identity morphism of SM and is only a morphism in $\mathfrak{Bund}$, not in $\mathfrak{VBund}_{\mathbb{R}}$, because it is not linear.

$\mathbf{C}^\infty \circ \mathbf{D}_0$ and $P_c : \mathbf{C}_0^\infty \circ \mathbf{D}_0^* \Rightarrow \mathbf{C}_0^\infty \circ \mathbf{D}_0^*$, $P_c : \mathbf{C}^\infty \circ \mathbf{D}_0^* \Rightarrow \mathbf{C}^\infty \circ \mathbf{D}_0^*$, which we denote by the same symbol. We then have by Proposition 3.7:

$$\begin{aligned} P \circ {}^c &= {}^c \circ P & P_c \circ {}^c &= {}^c \circ P_c, \\ P_c \circ {}^+ &= {}^+ \circ P & P \circ {}^+ &= {}^+ \circ P_c, \end{aligned} \quad (9)$$

i.e. if a spinor field u is a solution to the Dirac equation, then so are u^+ and u^c . (The adjoint and charge conjugation of u are defined pointwise.)

For a distribution v on D_0M we define the transpose P^* by $\langle P^*v, u \rangle := \langle v, Pu \rangle$ and similarly for P_c . In this way the transposes give rise to natural transformations $P^* : \mathbf{Distr} \circ \mathbf{D}_0 \Rightarrow \mathbf{Distr} \circ \mathbf{D}_0$ and $P_c^* : \mathbf{Distr} \circ \mathbf{D}_0^* \Rightarrow \mathbf{Distr} \circ \mathbf{D}_0^*$.

Lemma 3.9 *Let $\iota : \mathbf{C}^\infty \circ \mathbf{D}_0^* \Rightarrow \mathbf{Distr} \circ \mathbf{D}_0$ and $\iota : \mathbf{C}^\infty \circ \mathbf{D}_0 \Rightarrow \mathbf{Distr} \circ \mathbf{D}_0^*$ be the canonical natural transformations (see section 2.2 item 14). Then $P^* \circ \iota = \iota \circ P_c$ and $P_c^* \circ \iota = \iota \circ P$.*

Proof. This follows from the fact that for each object SM $\int_M \langle u, \nabla v \rangle d\text{vol}_g = - \int_M \langle \nabla u, v \rangle d\text{vol}_g$ if at least one of $u \in C^\infty(D_0M)$ and $v \in C^\infty(D_0^*M)$ is compactly supported. This in turn follows from $\langle \nabla v, u \rangle + \langle v, \nabla u \rangle = \nabla_a \langle v, \gamma^a u \rangle$ and Gauss' law. $\square$

One can find unique advanced and retarded fundamental solutions for the Dirac equation, both for spinors and cospinors [28, 14]:

Theorem 3.10 *There are unique natural transformations $S^\pm : \mathbf{C}_0^\infty \circ \mathbf{D}_0 \Rightarrow \mathbf{C}^\infty \circ \mathbf{D}_0$ and $S_c^\pm : \mathbf{C}_0^\infty \circ \mathbf{D}_0^* \Rightarrow \mathbf{C}^\infty \circ \mathbf{D}_0^*$ such that $S^\pm \circ P = P \circ S^\pm = \kappa$, $S_c^\pm \circ P_c = P_c \circ S_c^\pm = \kappa$ and such that for each $u \in C_0^\infty(D_0M)$, $v \in C_0^\infty(D_0^*M)$ we have $\text{supp}(S^\pm u) \subset J^\pm(\text{supp}(u))$, $\text{supp}(S_c^\pm u) \subset J^\pm(\text{supp}(u))$. Moreover,*

$$\begin{aligned} S^\pm \circ {}^c &= {}^c \circ S^\pm & S_c^\pm \circ {}^c &= {}^c \circ S_c^\pm, \\ S_c^\pm \circ {}^+ &= {}^+ \circ S^\pm & S^\pm \circ {}^+ &= {}^+ \circ S_c^\pm, \end{aligned}$$

$$\int \circ \langle, \rangle \circ (1 \otimes S^\pm) = \int \circ \langle, \rangle \circ (S_c^\mp \otimes 1).$$

Proof. The components of $S^\pm$ and $S_c^\pm$ are the advanced $(-)$ and retarded $(+)$ fundamental solutions for P and P_c , which are given by $S^\pm := (i\nabla + m)E^\pm$ and $S_c^\pm := (-i\nabla + m)E^\pm$ respectively, where $E^\pm$ are the unique advanced and retarded fundamental solutions for the normally hyperbolic operator $(i\nabla + m)(-i\nabla + m) = (-i\nabla + m)(i\nabla + m) = \nabla^2 + m^2$. We refer to [14] Theorem 2.1 for a detailed proof of the existence and uniqueness of these operators (see also [2] for the existence and uniqueness of $E^\pm$).

The naturality of $S^\pm$ and $S_c^\pm$ follows from their uniqueness and the naturality of P and P_c . In detail: for every morphism $\chi : SM_1 \rightarrow SM_2$ and every $f \in C_0^\infty(D_0M_1)$ the unique smooth solution to $Pu = \chi_* f$ on M_2 with $\text{supp}(u) \subset J^\pm(\text{supp}(\chi_* f))$ pulls back to a solution $v := \chi^* u$ of $Pv = f$ on M_1 with $\text{supp}(v) \subset J^\pm(\text{supp}(f))$. By uniqueness we must then have $u = S^\pm \chi_* f$ and $\chi^* u = S^\pm f$, i.e. $\chi^* \circ S^\pm \circ \chi_* = S^\pm$. The same holds for cospinors. The commutation of $S^\pm$ and $S_c^\pm$ with charge conjugation and adjoints follows from equation (9).

For arbitrary $u \in C_0^\infty(D_0M)$ and $v \in C_0^\infty(D_0^*M)$ we can find a $\phi \in C_0^\infty(M)$ which is identically 1 on the compact set $\text{supp}(S^\pm u) \cap \text{supp}(S_c^\mp v)$. We then compute:

$$\begin{aligned} \int_M \langle v, S^\pm u \rangle &= \int_M \langle P_c S_c^\mp v, \phi S^\pm u \rangle = \int_M \langle S_c^\mp v, P \phi S^\pm u \rangle \\ &= \int_M \langle S_c^\mp v, \phi P S^\pm u \rangle = \int_M \langle S_c^\mp v, u \rangle, \end{aligned}$$

which proves the last claim. $\square$

We define the advanced-minus-retarded fundamental solutions $S := S^- - S^+$ and $S_c := S_c^- - S_c^+$, which are natural transformations $S : \mathbf{C}_0^\infty \circ \mathbf{D}_0 \Rightarrow \mathbf{C}^\infty \circ \mathbf{D}_0$ and $S_c : \mathbf{C}_0^\infty \circ \mathbf{D}_0^* \Rightarrow \mathbf{C}^\infty \circ \mathbf{D}_0^*$ respectively.

3.4 The non-uniqueness of the functorial Dirac structure

We have seen that the (standard) structure of Dirac spinors and cospinors, adjoints, charge conjugation and the Dirac operator is entirely determined by the functor D_0 and the natural equivalences $^+, ^c$ and γ . We formalise this with a definition:

Definition 3.11 *By a Dirac structure $\mathcal{D} := (\mathbf{D}, ^+, ^c, \gamma)$ we mean a locally covariant spinor bundle $\mathbf{D}$ with a dual bundle $\mathbf{D}^*$, natural equivalences $^+ : \mathbf{D} \Leftrightarrow \mathbf{D}^*$, $^c : \mathbf{D} \Leftrightarrow \mathbf{D}$, and $^c : \mathbf{D}^* \Leftrightarrow \mathbf{D}^*$ in $\mathfrak{VBund}_{\mathbb{R}}$ and a natural transformation $\gamma : \mathbf{D} \Rightarrow \mathbf{T}^* \otimes \mathbf{D}$ in $\mathfrak{VBund}'_{\mathbb{C}}$, all of whose components cover the identity morphism and satisfying the relations (4) and $\langle \gamma_{SM}(v^+, v), n \rangle \geq 0$ for every time-like future pointing vector $n \in TM$.*

We call $\mathcal{D}_0 := (\mathbf{D}_0, ^+, ^c, \gamma)$ of Theorem 3.6 the standard Dirac structure.

The category $\mathfrak{DStruc}$ has all Dirac structures as objects and its morphisms $t : \mathcal{D}_1 \rightarrow \mathcal{D}_2$ are all natural transformations $t : \mathbf{D}_1 \Rightarrow \mathbf{D}_2$ whose components are injective morphisms covering the identity morphism and intertwining the adjoints, charge conjugation and γ as follows:

$$^{+2} \circ t = t \circ ^{+1}, \quad ^{c2} \circ t = t \circ ^{c1}, \quad \gamma_2 \circ (t \otimes t) = \gamma_1.$$

For each Dirac structure one can perform the constructions of subsection 3.3. Because the Dirac algebra D has a unique irreducible complex representation one might expect that the category $\mathfrak{DStruc}$ admits a corresponding unique initial object, perhaps up to isomorphism. This is an object from which there exists a morphism into any other object. However, as we will explain in this subsection there is a certain cohomological obstruction of the category $\mathfrak{SSpac}$ involved. We will first consider the standard Dirac structure, which would be a good candidate for an initial object, and prove the following weaker property:

Proposition 3.12 *Any morphism t from a Dirac structure $\mathcal{D}$ to the standard Dirac structure $\mathcal{D}_0$ is an isomorphism.*

Proof. Let $t : \mathcal{D} \rightarrow \mathcal{D}_0$ be a morphism. By the injectivity of the components of $t : \mathbf{D} \Rightarrow \mathbf{D}_0$ we see that the complex dimension of the fiber of DM is at most four. On the other hand, the vector bundles DM are modules for the Dirac algebra represented by γ . Because this algebra is simple, and because equations (4) exclude the trivial representation, we find that DM must have complex dimension at least four. Therefore, $t : \mathbf{D} \Rightarrow \mathbf{D}_0$ must be a natural equivalence and it follows that $t : \mathcal{D} \rightarrow \mathcal{D}_0$ is an isomorphism. $\square$

Corollary 3.13 *If we construct a Dirac structure $\mathcal{D}_\pi$ analogous to $\mathcal{D}_0$, but using a different representation π and matrices A, C , then $\mathcal{D}_\pi$ is isomorphic to $\mathcal{D}_0$.*

Proof. Because we use the same representation on all spacetimes we can construct a natural equivalence $t : \mathcal{D}_\pi \Leftrightarrow \mathcal{D}_0$ whose components are of the form $t_{SM}([E, z]) := [E, Lz]$ for some $L \in GL(4, \mathbb{C})$ which is independent of SM (cf. Theorem 3.5). $\square$

Corollary 3.14 *If $\mathcal{D} := (\mathbf{D}_0, ^{+1}, ^{c1}, \gamma')$ is any Dirac structure with the standard locally covariant Dirac spinor bundle $\mathbf{D}_0$, then $\mathcal{D}$ is isomorphic to the standard Dirac structure $\mathcal{D}_0$.*

Proof. At each point x in each object SM we can view γ'_a as matrices that represent the Dirac algebra in a representation π . Using the Fundamental Theorem 2.2 we write $\gamma'_a = L\gamma_a L^{-1}$ for some $L(x) \in GL(4, \mathbb{C})$. As γ'_a is well-defined on $\mathbf{D}_0$ we must have $\pi_0(S)\gamma'_a\pi_0(S^{-1}) = \gamma'_b\Lambda^b_a(S)$ for all $S \in \text{Spin}^0_{1,3}$. This also holds for the matrices γ , so we conclude from the Fundamental Theorem that $\pi_0(S)L(x) = c(x)L(x)\pi_0(S)$, where $c \equiv 1$ by taking $S = I$. We can now define a natural equivalence $t : \mathbf{D}_0 \Leftrightarrow \mathbf{D}_0$ by $[E, z] \mapsto [E, L(p(E))z]$ such that $\gamma' \circ t = t \circ \gamma$. If we also define $^{+2} := t \circ ^{+1} \circ t^{-1}$ and $^{c2} := t \circ ^{c1} \circ t^{-1}$, then $\mathcal{D} \Leftrightarrow (\mathbf{D}_0, ^{+2}, ^{c2}, \gamma) \Leftrightarrow \mathcal{D}_0$, where the last equivalence follows from the previous Corollary. $\square$

In fact, the proof of Corollary 3.13 shows that for any SM the quadruple $(DM, ^+, ^c, \gamma)$ is unique up to an isomorphism t_{SM} , if DM has four dimensional complex fibers. The isomorphism

t_{SM} itself, however, is only unique up to a sign. In other words, on each spin spacetime we find a discrete $\mathbb{Z}_2$ -symmetry that preserves all physical relations.¹⁶

Consider two Dirac structures $\mathcal{D}$ and $\mathcal{D}'$ whose locally covariant spinor bundles $\mathbf{D}$ and $\mathbf{D}'$ have four dimensional complex fibers. Comparing the action of these functors on morphisms of $\mathfrak{S}\mathfrak{P}\mathfrak{a}\mathfrak{c}$ one finds a diagram that commutes up to a sign. The existence of an initial object in the category $\mathfrak{D}\mathfrak{S}\mathfrak{t}\mathfrak{r}\mathfrak{u}\mathfrak{c}$ then boils down to the question whether one can choose signs for all spin spacetimes SM in such a way that all the diagrams commute. The answer is not at all obvious, but can be neatly formulated in terms of the first Stiefel-Whitney class of the category $\mathfrak{S}\mathfrak{P}\mathfrak{a}\mathfrak{c}$. To explain this we will briefly recall the definition of cohomology groups for categories (cf. [33]).

If $\mathfrak{C}$ is any category, we can first build a simplicial set from it called the *nerve* of the category (cf. [38]). A 0-simplex is simply an object of $\mathfrak{C}$, a 1-simplex is a morphism between two objects, a 2-simplex is a commutative triangle, etc. We will write Σ_n for the set of all n -simplices. For $n \geq 1$ every n -simplex has $n + 1$ *faces*, which are described by maps $\partial_j : \Sigma_n \rightarrow \Sigma_{n-1}$, $0 \leq j \leq n$, which remove the j^{th} vertex from the diagram.

To find the cohomology of $\mathfrak{C}$ with values in an Abelian group¹⁷ G , we define an n -cochain with values in G to be a map $v : \Sigma_n \rightarrow G$. We denote the set of n -cochains with values in G by $C^n(G)$ and we define the *coboundary* map $d : C^n(G) \rightarrow C^{n+1}(G)$ by

$$dv(s) := \sum_{j=0}^{n+1} (-1)^j v(\partial_j s), \quad s \in \Sigma_{n+1},$$

where we have written the group operation of G additively. One checks that $d^2 = 0$ and defines v to be *closed* iff $dv = 0$ and *exact* iff $v = dt$ for some $(n - 1)$ -cochain t . The sets of closed and exact n -cochains are denoted by $B^n(G)$ and $Z^n(G)$, respectively. They inherit an Abelian group structure from G and because $Z^n(G) \subset B^n(G)$ is necessarily normal one can define the j^{th} cohomology group as the quotient $H^n(G) := B^n(G)/Z^n(G)$.

Now let us return to the study of Dirac structures. Suppose that $\mathcal{D}$ and $\mathcal{D}'$ both have four dimensional complex fibers. Without loss of generality we may assume that both Dirac structures coincide on each spin spacetime, but the action of their locally covariant spinor bundles on a morphism χ agrees only up to a sign $v(\chi) \in \{\pm 1\}$. We can view $v : \chi \mapsto v(\chi)$ as a 1-cochain on the category $\mathfrak{S}\mathfrak{P}\mathfrak{a}\mathfrak{c}$ with values in $\mathbb{Z}_2 = \{0, 1\}$, where 0 corresponds to $+1$ and 1 to -1 . Notice that for a composition of morphisms $\chi = \chi_1 \circ \chi_2$ we find $v(\chi) = v(\chi_1) + v(\chi_2)$ in $\mathbb{Z}_2$, because the Dirac structures are both functorial. In cohomological terms this means precisely that $dv = 0$.

If there is a natural equivalence $t : \mathcal{D} \Leftrightarrow \mathcal{D}'$, then the components t_{SM} are automorphisms of the Dirac structure at each SM , i.e. $t_{SM} = \pm 1$, that compensate for all the minus signs in v . If we view t as a 0-cochain with values in $\mathbb{Z}_2$, this means exactly that $v = dt$. So we have proved:

Theorem 3.15 *The number of inequivalent Dirac structures whose locally covariant spinor bundles have four dimensional complex fibers equals the number of first Stiefel-Whitney classes of the category $\mathfrak{S}\mathfrak{P}\mathfrak{a}\mathfrak{c}$, i.e. the number of elements in $H^1(\mathbb{Z}_2)$.*

Remark 3.16 *For scalar and vector fields the problem above can be avoided in a natural way. Taking $\mathcal{L}_+^\dagger$ in the defining (four-vector) representation, the vector bundle associated to $F_+^\dagger M$ is just the tangent bundle TM . A morphism in $\mathfrak{S}\mathfrak{P}\mathfrak{a}\mathfrak{c}$ determines a unique morphism on the tangent bundle, so no topological obstructions occur. Similarly for the scalar field, where one uses the trivial one dimensional representation of $\mathcal{L}_+^\dagger$, whose associated vector bundle is $\Lambda^0(M) = M \times \mathbb{R}$. Again a morphism in $\mathfrak{M}\mathfrak{a}\mathfrak{n}^n$ automatically determines a unique morphism on these associated vector bundles, now by the requirement that the volume element is preserved.*

In general one is dealing with representations of $\text{Spin}_{1,3}^0$ and associates to each morphism in $\mathfrak{S}\mathfrak{P}\mathfrak{a}\mathfrak{c}$ an intertwining operator between such representations. For the associated vector bundles of SM , the physical requirements that we imposed on the bundle morphisms, concerning the adjoint

¹⁶This may be compared to [6], who use complex spinor structures and then find a local (gauge) symmetry instead of our more restricted global symmetries.

¹⁷[33] also considers the non-Abelian case, which is much more involved.

and charge conjugation maps and γ , reduce the intertwiners exactly to a choice of lifting $\mathcal{L}_+^\dagger$ to its double cover. In this way it leads to the same first Stiefel-Whitney class that characterises the number of spin structures on a manifold. For the general case it is expected that one needs a non-Abelian cohomology theory to quantify the obstruction for finding initial objects.

4 The locally covariant quantum Dirac field

After our discussion of the classical Dirac field in section 3 we now turn to the quantum Dirac field, its construction, its Hadamard states and its relative Cauchy evolution.

4.1 Quantisation of the free Dirac field

First we will quantise the free Dirac field in a generally covariant way and establish some of its properties. For this purpose we also present the main ideas of locally covariant quantum field theory as introduced in [4] (see also [16]).

In the following any quantum physical system will be described by a topological $*$ -algebra $\mathcal{A}$ with a unit I , whose self-adjoint elements are the observables of the system. An injective and continuous $*$ -homomorphism expresses the notion of a subsystem, whereas a state is described by a normalised and positive continuous linear functional ω , i.e. $\omega(A^*A) \geq 0$ for all $A \in \mathcal{A}$ and $\omega(I) = 1$. The state space of $\mathcal{A}$ is the set of all states and is denoted by $\mathcal{A}_1^{*+}$. Every state gives rise to a GNS-representation π_ω (see [37] Theorem 8.6.2.), which is characterised uniquely, up to unitary equivalence, by the GNS-quadruple $(\pi_\omega, \mathcal{H}_\omega, \Omega_\omega, \mathcal{D}_\omega)$. Here $\mathcal{H}_\omega$ is the Hilbert space on which $\pi_\omega(\mathcal{A})$ acts as (possibly unbounded) operators with the dense, invariant domain $\mathcal{D}_\omega := \pi_\omega(\mathcal{A})\Omega_\omega$. The vector Ω_ω is cyclic and satisfies $\omega(A) = \langle \Omega_\omega, \pi_\omega(A)\Omega_\omega \rangle$ for all $A \in \mathcal{A}$.

The collection of all systems forms a category $\mathfrak{TAlg}$:

Definition 4.1 *The category $\mathfrak{TAlg}$ has as its objects all unital topological $*$ -algebras $\mathcal{A}$ and as its morphisms all continuous and injective $*$ -homomorphisms α such that $\alpha(I) = I$.*

A locally covariant quantum field theory is a (covariant) functor $\mathbf{A} : \mathfrak{Spc} \rightarrow \mathfrak{TAlg}$, written as $SM \mapsto \mathcal{A}_{SM}$, $\chi \mapsto \alpha_\chi$.

A locally covariant quantum field theory $\mathbf{A}$ is called causal if and only if any pair of morphisms $\psi_i : SM_i \rightarrow SM$, $i = 1, 2$, such that $\psi_1(\mathcal{M}_1) \subset (\psi_2(\mathcal{M}_2))^\perp$ in $\mathcal{M}$ yields $[\alpha_{\psi_1}(\mathcal{A}_{SM_1}), \alpha_{\psi_2}(\mathcal{A}_{SM_2})] = \{0\}$ in $\mathcal{A}_{SM}$.

A locally covariant quantum field theory $\mathbf{A}$ satisfies the time-slice axiom iff for all morphisms $\psi : SM_1 \rightarrow SM_2$ such that $\psi(\mathcal{M}_1)$ contains a Cauchy surface for $\mathcal{M}_2$ we have $\alpha_\psi(\mathcal{A}_{SM_1}) = \mathcal{A}_{SM_2}$.

Notice that the condition $\psi_1(\mathcal{M}_1) \subset (\psi_2(\mathcal{M}_2))^\perp$ is symmetric in $i = 1, 2$. The causality condition formulates how the quantum physical system interplays with the classical gravitational background field, whereas the time-slice axiom expresses the existence of a causal dynamical law.

We now fix a choice of Dirac structure $\mathcal{D} := (\mathbf{D}, ^+, ^c, \gamma)$, in order to turn the free Dirac field into a locally covariant field theory. Because we want to impose the canonical anti-commutation relations it will also be convenient to quantise spinor and cospinor fields simultaneously by introducing the following terminology:

Definition 4.2 *The locally covariant double spinor bundle is the covariant functor $D \oplus D^*$. We define the following natural equivalences and natural transformations on this bundle, indicated by their components at SM :*

$$\begin{aligned} (p \oplus q)^c &:= p^c \oplus q^c & (p \oplus q)^+ &:= q^+ \oplus p^+ \\ \gamma_\mu(p \oplus q) &:= (\gamma_\mu p) \oplus (\gamma_\mu q) & \langle p \oplus q, p' \oplus q' \rangle &:= \langle p^+, p' \rangle + \langle q', q^+ \rangle \\ R(p \oplus q) &:= p \oplus (-q). \end{aligned}$$

A double spinor (field) is an element of $C^\infty(DM \oplus D^*M)$. A double test-spinor (field) is an element of $C_0^\infty(DM \oplus D^*M)$. The adjoint, charge conjugation and other operations are defined

pointwise. We also define the operator $P := P \oplus P_c$, its advanced $(-)$ and retarded $(+)$ fundamental solutions $S^\pm(u \oplus v) := (S^\pm u) \oplus (S_c^\pm v)$ and $S := S^- - S^+$.

The exterior tensor product $\mathcal{V}_1 \boxtimes \mathcal{V}_2$ of two vector bundles $\mathcal{V}_i$ with fiber V_i over manifolds $\mathcal{M}_i$, $i = 1, 2$, is the vector bundle over $\mathcal{M}_1 \times \mathcal{M}_2$ whose fiber is $V_1 \otimes V_2$ and whose local trivialisations are determined by $(O_1 \times O_2) \times (V_1 \otimes V_2)$, where $O_i \times V_i$ are local trivialisations of $\mathcal{V}_i$.

The Dirac Borchers-Uhlmann algebra $\mathcal{F}_{SM}^0$ on a spin spacetime SM is the topological $*$ -algebra

$$\mathcal{F}_{SM}^0 := \oplus_{n=0}^{\infty} C_0^\infty((DM \oplus D^*M)^{\boxtimes n}),$$

where the direct sum is algebraic (i.e. only finitely many non-zero summands are allowed) and

1. the product is given by continuous linear extension of $f_1 \cdot f_2 := f_1 \boxtimes f_2$,
2. the $*$ -operation is given by continuous antilinear extension of

$$(f_1 \boxtimes \dots \boxtimes f_n)^* := f_n^+ \boxtimes \dots \boxtimes f_1^+,$$

3. as a topological vector space $\mathcal{F}_{SM}^0$ is the strict inductive limit $\mathcal{F}_{SM}^0 = \cup_{N=0}^{\infty} \oplus_{n=0}^N C_0^\infty((DM \oplus D^*M)^{\boxtimes n}|_{K_N^{\times n}})$, where K_N is an exhausting and increasing sequence of compact subsets of $\mathcal{M}$ and the test-section space of the restricted vector bundle $(DM \oplus D^*M)^{\boxtimes n}|_{K_N^{\times n}}$ is given the test-section topology.

The topology of $\mathcal{F}_{SM}^0$ is such that a state is given by a sequence of n -point distributional sections ω_n of $(DM \oplus D^*M)^{\boxtimes n}$. A morphism $\chi: SM_1 \rightarrow SM_2$ in $\mathfrak{S}\mathfrak{P}\mathfrak{a}\mathfrak{c}$ determines a unique morphism $\alpha_\chi: \mathcal{F}_{SM_1}^0 \rightarrow \mathcal{F}_{SM_2}^0$ that is given by the algebraic and continuous extension of the morphism $DM_1 \oplus D^*M_1 \rightarrow DM_2 \oplus D^*M_2$ that is supplied by the functor $\mathbf{D}$. Together with this map on morphisms the map $SM \mapsto \mathcal{F}_{SM}^0$ becomes a locally covariant quantum field theory $\mathbf{F}^0: \mathfrak{S}\mathfrak{P}\mathfrak{a}\mathfrak{c} \rightarrow \mathfrak{T}\mathfrak{A}\mathfrak{I}\mathfrak{g}$. Our next task will be to divide out the ideals that generate the dynamics and the canonical anti-commutation relations.

We define the natural transformation $(,): (\mathbf{C}_0^\infty \circ (\mathbf{D} \oplus \mathbf{D}^*)) \otimes_{\mathbb{R}} (\mathbf{C}_0^\infty \circ (\mathbf{D} \oplus \mathbf{D}^*)) \Rightarrow \mathbb{C}$ whose components are the sesquilinear forms:

$$(f_1, f_2) := i \int_M \langle f_1, RS f_2 \rangle.$$

Note that this is indeed a natural transformation, because it can be written as a composition of natural transformations including $\int$, $\langle, \rangle$, $+$ and κ .

Lemma 4.3 *On each object SM the sesquilinear form $(,)$ is Hermitean, $\overline{(f_1, f_2)} = (f_1^c, f_2^c) = (f_2, f_1)$, and there holds $(f_1^+, f_2^+) = (f_2, f_1)$. For any spacelike Cauchy surface $C \subset M$ with future pointing unit normal vector field n^a we have*

$$(u_1 \oplus v_1, u_2 \oplus v_2) = \int_C \langle (Su_1)^+, \not{n}(Su_2) \rangle + \langle S_c v_2, \not{n}(S_c v_1)^+ \rangle. \quad (10)$$

Proof. The symmetry properties follow straightforwardly from the computational rules of Theorems 3.6 and 3.10. For the last statement we also need a partial integration (see e.g. [42] equation (B.2.26) for Gauss' law) and we use the Dirac equation:

$$\begin{aligned} & (u_1 \oplus v_1, u_2 \oplus v_2) \\ &= i \int_{J^+(C)} \langle P_c S_c^- u_1^+, Su_2 \rangle + \langle P_c S_c^- v_2, S v_1^+ \rangle + i \int_{J^-(C)} \langle P_c S_c^+ u_1^+, Su_2 \rangle + \langle P_c S_c^+ v_2, S v_1^+ \rangle \\ &= - \int_{J^+(C)} \nabla_a \langle S_c^- u_1^+, \gamma^a Su_2 \rangle + \nabla_a \langle S_c^- v_2, \gamma^a S v_1^+ \rangle \\ &\quad - \int_{J^-(C)} \nabla_a \langle S_c^+ u_1^+, \gamma^a Su_2 \rangle + \nabla_a \langle S_c^+ v_2, \gamma^a S v_1^+ \rangle \end{aligned}$$

$$\begin{aligned}
&= \int_C n_a \langle S_c^- u_1^+, \gamma^a S u_2 \rangle + n_a \langle S_c^- v_2, \gamma^a S v_1^+ \rangle - \int_C n_a \langle S_c^+ u_1^+, \gamma^a S u_2 \rangle + n_a \langle S_c^+ v_2, \gamma^a S v_1^+ \rangle \\
&= \int_C \langle (S u_1)^+, \not{S} (S u_2) \rangle + \langle S_c v_2, \not{S} (S_c v_1)^+ \rangle.
\end{aligned}$$

□

From equation (10) we notice that $(\cdot, \cdot)$ is positive semi-definite and hence defines a (degenerate) inner product. We proceed by dividing $\mathcal{F}_{SM}^0$ by the closed ideal J_{SM} of $\mathcal{F}_{SM}^0$ generated by all elements of the form Pf or $f_1^+ \cdot f_2 + f_2 \cdot f_1^+ - (f_1, f_2)I$.

Theorem 4.4 *The ideal J_{SM} is a $*$ -ideal and for any morphism $\chi : SM_1 \rightarrow SM_2$ we have $\alpha_\chi(J_{SM_1}) \subset J_{SM_2}$. We can define the locally covariant quantum field theory $\mathbf{F} : \mathfrak{Spac} \rightarrow \mathfrak{Alg}$ which assigns to every spin spacetime SM the C^* -algebra $\mathcal{F}_{SM} := \mathcal{F}_{SM}^0 / J_{SM}$.*

Proof. The elements that generate J_{SM} are invariant under adjoints and under a morphism they are mapped to elements of the same form. This proves the first statement. It follows that the quotients $\mathcal{F}_{SM}^0 / J_{SM}$ are topological $*$ -algebras and that a morphism $\alpha_\chi : \mathcal{F}_{SM_1}^0 \rightarrow \mathcal{F}_{SM_2}^0$ descends to the quotients as a well-defined morphism. That each algebra $\mathcal{F}_{SM}^0 / J_{SM}$ has a C^* -norm follows from the fact that they are the inductive limits of finite dimensional Clifford algebras ([1]). The morphisms on the quotients are necessarily continuous in the norm and therefore extend to morphisms on the C^* -algebras $\mathcal{F}_{SM}$. □

Definition 4.5 *A locally covariant quantum field in the locally covariant vector bundle $\mathbf{V}$ for the locally covariant quantum field theory $\mathbf{A}$ is a natural transformation $\Phi : \mathbf{C}_0^\infty \circ \mathbf{V}^* \Rightarrow \mathbf{f} \circ \mathbf{A}$, where we let $\mathbf{f} : \mathfrak{Alg} \rightarrow \mathfrak{Vec}$ be the forgetful functor.*

We define the locally covariant quantum fields $B : \mathbf{D} \otimes \mathbf{D}^ \Rightarrow \mathbf{F}$, $\psi : \mathbf{D}^* \Rightarrow \mathbf{F}$ and $\psi^+ : \mathbf{D} \Rightarrow \mathbf{F}$ by $B_{SM}(f) := 0 \oplus f \oplus 0 \oplus \dots + J_{SM}$, $\psi_{SM}(v) := B_{SM}(0 \oplus v)$ and $\psi_{SM}^+(u) := B_{SM}(u \oplus 0)$.*

That the latter really are locally covariant quantum fields is a consequence of

Proposition 4.6 *The operator-valued maps $B_{SM}, \psi_{SM}, \psi_{SM}^+$ are C^* -algebra-valued distributions and:*

1. $P \circ \psi = 0$ and $P_c \circ \psi^+ = 0$,
2. $\psi_{SM}^+(u) = \psi_{SM}(u^+)^*$,
3. $\{\psi_{SM}^+(u), \psi_{SM}(v)\} = (v^+ \oplus 0, u \oplus 0)I = -i \int_M \langle v, S u \rangle I$ and the other anti-commutators vanish.

Proof. The first item is $P B_{SM}(f) = B_{SM}(P^* f) = B_{SM}(P f) = 0$, where P^* is the formal adjoint of P . The last two items follow from the definitions of ψ_{SM} and ψ_{SM}^+ and the properties of B_{SM} after a straight-forward computation.

It remains to show that ψ_{SM}, ψ_{SM}^+ are C^* -algebra-valued distributions, because the result for B_{SM} then follows. The C^* -subalgebra of $\mathcal{F}_{SM}$ generated by $I, \psi_{SM}(v), \psi(v)_{SM}^*$ is a Clifford algebra which is isomorphic to $M(2, \mathbb{C})$ and an explicit isomorphism is given by $\psi_{SM}(v) \mapsto \begin{pmatrix} 0 & \sqrt{c} \\ 0 & 0 \end{pmatrix}$, where $c = (0 \oplus v, 0 \oplus v) = -i \int_M \langle v, S v^+ \rangle > 0$. It follows that $\|\psi_{SM}(v)\| = \sqrt{c}$ is the operator norm of the corresponding matrix, i.e.¹⁸

$$\|\psi_{SM}(v)\|^2 = -i \int_M \langle v, S v^+ \rangle d\text{vol}_g.$$

In the test-spinor topology we then have continuous maps $v \mapsto v \oplus v^+ \mapsto -i \int_M \langle v, S v^+ \rangle$, from which it follows that $v \mapsto \psi_{SM}(v)$ is norm continuous, i.e. it is a C^* -algebra-valued distribution. The proof for ψ_{SM}^+ is analogous. □

¹⁸The factor 2 in [17] Remark 2, p.340 seems to be erroneous.

Note that the last two conditions of Proposition 4.6 can also be formulated in terms of natural transformations, because the algebraic operations in $\mathcal{F}_{SM}$ can be expressed as such. The theory $\mathbf{F}$ is the quantised free Dirac field and ψ (ψ^+) is the locally covariant Dirac (co)spinor field. Alternatively we could have used the algebras $\mathcal{F}_{SM}^0/J_{SM}$ themselves instead of completing them to C^* -algebras.

To see that the anti-commutator is the canonical one (cf. [28]) we apply Proposition 2.4c) of [14], which says that $S|_{C \times C} = -i\delta\rlap{/\kern-0.1em}/$ for a Cauchy surface C with future pointing normal vector field n . Comparing with equation (8) and using $\rlap{/\kern-0.1em}/^2 = I$ we then find

$$\{-i\psi_{SM}^+(\rlap{/\kern-0.1em}/(x)), \psi_{SM}(y)\} = - \int_M \langle y, S\rlap{/\kern-0.1em}/x \rangle I = i\delta(y, x)I$$

as expected.

So far our construction depends on the choice of a Dirac structure, although naturally equivalent Dirac structures yield naturally equivalent theories and quantum fields. The following theorem restricts attention to the observable algebra, dividing out the freedom of choice completely and yielding a unique theory, but for many purposes it is not convenient to use it directly because it lacks locally covariant Dirac (co)spinor fields.

Theorem 4.7 *Let $\mathbf{B} : \mathfrak{S}\mathfrak{P}\mathfrak{a}\mathfrak{c} \rightarrow \mathfrak{T}\mathfrak{A}\mathfrak{l}\mathfrak{g}$ be the locally covariant quantum field theory that assigns to each spin spacetime SM the C^* -subalgebra of $\mathcal{F}_{SM}$ generated by all even polynomials in elements $B(f)$, with the induced action on morphisms. For all Dirac structures with four dimensional complex fibers the resulting theories $\mathbf{B}$ are isomorphic.*

Proof. The algebras $\mathcal{B}_{SM}$ generated by the even polynomials are C^* -algebras. Morphisms respect evenness and so restrict to morphisms on $\mathcal{B}$, making $\mathbf{B}$ a well-defined locally covariant quantum field theory. Now consider two Dirac structures $\mathcal{D}$ and $\mathcal{D}_0$ with associated functors $\mathbf{F}, \mathbf{B}$ and $\mathbf{F}_0, \mathbf{B}_0$. If both Dirac structures have four dimensional complex fibers, then we infer from the comment below Corollary 3.13 that there are $*$ -isomorphisms $\alpha_{SM} : \mathcal{F}_{SM} \rightarrow (\mathcal{F}_0)_{SM}$ such that for any morphism $\chi : SM_1 \rightarrow SM_2$ we have $\alpha_{SM_2} \circ \alpha_\chi = \epsilon_\chi \cdot (\alpha_0)_\chi \circ \alpha_{SM_1}$, where $\epsilon_\chi = \pm 1$ depends only on χ . It follows from the evenness that the α_{SM} descend to $*$ -isomorphisms $\alpha_{SM} : \mathcal{B}_{SM} \rightarrow (\mathcal{B}_0)_{SM}$ that intertwine with the morphisms. Hence, $\mathbf{B}$ and $\mathbf{B}_0$ are naturally equivalent. $\square$

Proposition 4.8 *The locally covariant quantum field theory $\mathbf{B} : \mathfrak{S}\mathfrak{P}\mathfrak{a}\mathfrak{c} \rightarrow \mathfrak{T}\mathfrak{A}\mathfrak{l}\mathfrak{g}$ of Theorem 4.7 is causal and satisfies the time-slice axiom.*

Proof. Causality follows from the anti-commutation relations,

$$\begin{aligned} & [B_{SM}(f_1)B_{SM}(f_2), B_{SM}(f_3)] \\ &= B_{SM}(f_1) \{B_{SM}(f_2), B_{SM}(f_3)\} - \{B_{SM}(f_1), B_{SM}(f_3)\} B_{SM}(f_2) \\ &= (f_2, f_3)B_{SM}(f_1) - (f_1, f_3)B_{SM}(f_2), \end{aligned}$$

together with the support properties of S . For the time-slice axiom we let $\chi : SM \rightarrow SM'$ be a morphism in $\mathfrak{S}\mathfrak{P}\mathfrak{a}\mathfrak{c}$, covering a morphism $\psi : M \rightarrow M'$ in $\mathfrak{S}\mathfrak{P}\mathfrak{a}\mathfrak{c}$, such that $N := \psi(M) \subset M'$ contains a Cauchy surface $C \subset M'$. Then we can choose Cauchy surfaces $C^\pm \subset N$ such that $C^\pm \subset I^\pm(C)$ and a smooth partition of unity ϕ^+, ϕ^- with $\text{supp } \phi^\pm \subset J^\pm(C^\mp)$. Let $f \in C_0^\infty(DM \oplus D^*M)$ and write

$$f = P(S^+f - \phi^+Sf) + \tilde{f}, \quad (11)$$

where $\tilde{f} := P(\phi^+Sf) = -P(\phi^-Sf)$ is supported in $J^+(C^-) \cap J^-(C^+) \subset N$ and $\phi^+Sf - S^+f$ has compact support. Hence, $B_{SM_2}(f) = B_{SM_2}(\tilde{f}) = \alpha_\chi(B_{SM_1}(\chi^*(\tilde{f})))$. Because the algebra $\mathcal{F}_{M'}$ is generated by such elements this shows that α_χ is a $*$ -isomorphism. $\square$

Remark 4.9 *A Majorana spinor is a spinor u such that $u = u^c$. In this case the adjoint is anti-Majorana: $u^{+c} = -u^{c+} = -u^+$. We call a double spinor $f = u \oplus v$ Majorana iff u and v^+ are*

Majorana, which means that $f^c = Rf$. Such spinors are sections of a subbundle of the Dirac spinor bundle, which can be described by a Majorana representation. Notice that every spinor is a unique complex linear combination of Majorana spinors.

To quantise Majorana spinors we note that $\langle h^c, f \rangle = \langle h^+, f^{c+} \rangle$. This leads us to define the charge conjugation on the quantised fields¹⁹ by $\psi^c(v) := \psi^+(v^{c+})$ and $\psi^{+c}(u) := \psi(u^{c+})$, or equivalently $B^c(f) := B(f^{c+}) = B(f^c)^*$. We impose the Majorana condition $B^c(f) = B(Rf)$ by dividing out the ideal generated by all elements of the form $B(f - Rf^{c+})$. More precisely, if $\mathcal{H}$ is the Hilbert space obtained from $C_0^\infty(DM \oplus D^*M)$ by dividing out the ideal of double spinors f for which $(f, f) = 0$, then there is an orthogonal decomposition $\mathcal{H} = \mathcal{H}_+ \oplus \mathcal{H}_-$, where the elements in $\mathcal{H}_\pm$ satisfy $Rf^{c+} = \pm f$. Indeed, every double spinor can be written as $f = f_+ + if_-$, where $f_\pm := \frac{1}{2}(f \pm Rf^{c+})$ are in $\mathcal{H}_\pm$ and the orthogonality follows from Lemma 4.3. For the C^* -algebraic quantisation we then have $\mathcal{F} = \mathcal{F}_+ \otimes \mathcal{F}_-$, where $\mathcal{F}_-$ is the C^* -algebra of quantised Majorana spinors and $\mathcal{F}_+$ the C^* -algebra of quantised anti-Majorana spinors (see [3] section 5.2). The generators $\psi(v)$ and $\psi^+(u)$ of $\mathcal{F}_-$ satisfy the additional relation $\psi^c = \psi$ and $\psi^{+c} = -\psi^+$.

4.2 Hadamard states

After Radzikowski's result [32] that a for a scalar field state is of Hadamard form if and only if its wave front set has a certain form, several people set out to extend this result to the Dirac field, or more general quantum fields [26, 22, 34]. All three papers have provided an original contribution in their method of proof, but upon careful analysis they all have minor gaps. We feel that it is justified to comment on this here and to provide the necessary results to fill any remaining gaps.

The most general results are the most recent ones, due to Sahlmann and Verch [34], who set out to prove the equivalence of the Hadamard form of a state, defined in terms of the Hadamard parametrix, with a wave front set condition analogous to the scalar field case. One of the techniques used is the scaling limit, but the proof of their Proposition 2.8, which relates the wave front set of a distribution to that of its scaling limit, is in our opinion insufficient (see the footnote on page 28). In the appendix we prove a similar statement as Proposition A.2, thereby filling any gap in [34] and establishing the desired equivalence on a firm ground. For the Dirac field, Hollands has proved that this wave front set condition implies a specific form of the polarisation set ([23] Theorem 4.1).

The scaling limit result can also be used to find the wave front sets of the advanced and retarded fundamental solutions $E^\pm$ of normally hyperbolic operators on a globally hyperbolic spacetime, a result that we prove as Theorem A.5. Our proof is largely analogous to the work of Radzikowski and the outcome is in direct analogy to the results of Duistermaat and Hörmander [15] for the scalar case. To find the wave front sets of the fundamental solutions $S^\pm$ for the Dirac equation we use (and correct) an idea of [23].

Finally we comment on the results by Kratzert [26], which use a spacetime deformation argument to compute the wave front set and polarisation set of Hadamard states. This result has a gap, already identified in [34], concerning the case of points $(x, \xi; y, \xi')$ where either $\xi = 0$ or $\xi' = 0$, which prevents the propagation of the singularity from the original to the deformed spacetime. This gap can be avoided using either a propagation of Hadamard form result as in [34], or using the commutation or anti-commutation relations and the explicit form of $WF(E)$, respectively $WF(S)$. The latter argument, which appears to be implicit in Radzikowski's paper [32], works as follows: when $(x, \xi; y, 0) \in WF(\omega_2)$ then also $(y, 0; x, \xi) \in WF(\omega_2)$ by the (anti-)commutation relations and the fact that $WF(E)$ (or $WF(S)$) has no points with either entry equal to 0. Using the calculus of Hilbert-space-valued distributions, Theorem A.4, we then find that both $(x, \xi; x, -\xi) \in WF(\omega_2)$ and $(x, -\xi; x, \xi) \in WF(\omega_2)$. Because $\xi \neq 0$ (by definition the wave front set does not contain the zero covector) these points can both be propagated into a deformed spacetime, where $WF(\omega)$ is known to satisfy the required microlocal condition. This, however, leads to a contradiction, because $WF(\omega_2) \cap -WF(\omega_2) = \emptyset$ and hence $\xi = 0$. Therefore, $WF(\omega_2)$ cannot contain points with one of the covectors equal to 0.

¹⁹Our definition differs slightly from that of [11].

After these historical notes we feel free to define the notion of Hadamard states directly in terms of a wave front set condition, rather than using the Hadamard parametrix. If ω is a state on $\mathcal{F}_{SM}$ then we may consider the GNS-representation $(\mathcal{H}_\omega, \pi_\omega, \Omega_\omega)$ associated to ω and the $\mathcal{H}_\omega$ -valued distribution on $DM \oplus D^*M$ defined by:

$$v_\omega(f) := \pi_\omega(B_{SM}(f))\Omega_\omega.$$

Definition 4.10 *A state ω on $\mathcal{F}_{SM}$ is called Hadamard if and only if*

$$WF(v_\omega) = \mathcal{N}^+ := \{(x, \xi) \in T^*M \mid \xi^2 = 0, \xi^\mu \text{ is future pointing or } 0\}.$$

A state ω on $\mathcal{B}_{SM}$ is called Hadamard if and only if it can be extended to a Hadamard state on $\mathcal{F}_{SM}$. The set of all Hadamard states on $\mathcal{B}_{SM}$ will be denoted by $\mathcal{S}_{SM}$.

Note that every state on $\mathcal{B}_{SM}$ can be extended to $\mathcal{F}_{SM}$, by the Hahn-Banach Theorem and Proposition 4.6. The Hadamard condition is independent of the choice of extension, because it depends solely on the two-point distribution as the following proposition shows (cf. [34], we give a short proof using the more advanced microlocal techniques developed in the appendix).

Proposition 4.11 *For a state ω on $\mathcal{F}_{SM}$ the following conditions are equivalent:*

1. ω is Hadamard,
2. $WF(v_\omega) \subset \mathcal{N}^+$,
3. the two-point distribution $\omega_2(f_1, f_2) := \omega(B_{SM}(f_1)B_{SM}(f_2))$ has

$$WF(\omega_2) = \mathcal{C} := \{(x, -\xi; y, \xi') \in T^*M^{\times 2} \setminus \mathcal{Z} \mid (x, \xi) \sim (y, \xi'), (x, \xi) \in \mathcal{N}^+\},$$

where $(x, \xi) \sim (y, \xi')$ if and only if there is an affinely parameterised light-like geodesic from x to y to which ξ, ξ' are cotangent,

4. *there is a two-point distribution w such that $\omega_2(f_1, f_2) = iw(Pf_1, f_2)$ and $WF(w) = \mathcal{C}$.*

Proof. First note that ω_2 is a bidistribution on $DM \oplus D^*M$, because B_{SM} is an $\mathcal{F}_{SM}$ -valued distribution and multiplication in $\mathcal{F}_{SM}$ and ω are continuous. By Theorem A.4 the third statement implies the first, which trivially implies the second. To show that the second statement implies the third we use the argument of [39], Proposition 6.1. By Theorem A.4 we see that $WF(\omega_2) \subset \mathcal{N}^- \times \mathcal{N}^+$, where $\mathcal{N}^- := -\mathcal{N}^+$. Defining $\tilde{\omega}_2(f_1, f_2) := \omega_2(f_2, f_1)$ we find $WF(\tilde{\omega}_2) \cap WF(\omega_2) = \emptyset$. Now, $(\omega_2 + \tilde{\omega}_2)(f_1, f_2) = i \int_M \langle f_1, RSf_2 \rangle$, so $WF(\omega_2) \cup WF(\tilde{\omega}_2) = WF(S) = WF(E)$ by Proposition A.7 and hence $WF(\omega_2) = WF(E) \cap \mathcal{N}^- \times \mathcal{N}^+ = \mathcal{C}$ by Corollary A.6.

Now, assume that $\omega_2(f_1, f_2) = iw(Pf_1, f_2)$, where $WF(w) = \mathcal{C}$. Then $WF(\omega_2) = WF((P^* \otimes I)w) \subset WF(w) = \mathcal{C}$. It follows that $WF(v_\omega) \subset \mathcal{N}^+$. For the converse we suppose that ω is Hadamard and we choose a smooth real-valued function $\phi^\pm$ on M such that $\phi^+ \equiv 0$ to the past of some Cauchy surface C_- and such that $\phi^- := 1 - \phi^+ \equiv 0$ to the future of another Cauchy surface C_+ . We then define $w(f_1, f_2) := -i\omega_2(\phi^+ S^- f_1 + \phi^- S^+ f_1, f_2)$. Note that w is a bidistribution which is well-defined, because $\phi^+ S^- f_1$ and $\phi^- S^+ f_1$ are compactly supported. By construction $iw(Pf_1, f_2) = \omega_2(f_1, f_2)$. We now estimate the wave front set of w as follows. The wave front sets of $S^\pm$ are determined in Proposition A.7. Then we may apply Theorem 8.2.9 and 8.2.13 in [24] (in combination with equation (17)) to estimate the wave front sets of the tensor products $\phi^\pm(x)S^\mp(x, y)\delta(x', y')$ and the compositions in $iw(x, x') = \sum_\pm \int \omega_2(y, y')(\phi^\pm(x)S^\mp(x, y)\delta(x', y'))$ respectively and, using $WF(\omega_2) = \mathcal{C}$, we find:

$$WF(iw) \subset \cup_\pm WF(S^\mp \otimes \delta) \circ WF(\omega_2) \subset WF(\omega_2) = WF((P^* \otimes I)w) \subset WF(w),$$

i.e. $WF(w) = WF(\omega_2) = \mathcal{C}$. □

The second characterisation in Proposition 4.11 is especially useful, because it shows we do not need to compute the entire wave front set, as long as we can estimate it. Employing similar techniques as above one can use the anti-commutation relations and the wave front set of ω_2 to estimate the wave front sets of all higher n -point distributions [35], showing that a Hadamard state necessarily satisfies the microlocal spectrum condition (μ SC) of [5] and it follows that the set of such states is closed under operations from the algebra. We formulate this and other properties of Hadamard states in the following

Proposition 4.12 *The set $\mathcal{S}_{SM}$ of all Hadamard states on $\mathcal{B}_{SM}$ satisfies:*

1. $\alpha_\chi^*(\mathcal{S}_{SM_1}) \subset \mathcal{S}_{SM_2}$ for every morphism $\chi: SM_1 \rightarrow SM_2$,
2. $\mathcal{S}_{SM}$ is closed under operations from $\mathcal{B}_{SM}$,
3. $\alpha_\chi^*(\mathcal{S}_{SM_1}) = \mathcal{S}_{SM_2}$ for every morphism $\chi: SM_1 \rightarrow SM_2$ such that $\psi(\mathcal{M}_1)$ contains a Cauchy surface of $\mathcal{M}_2$.

Proof. The first property follows from Theorem 4.11 and the fact that wave front sets are local and geometric objects (cf. [24] Ch. 8). The second property relies on the anti-commutation relations, which implies that the truncated n -point distributions are totally anti-symmetric (cf. [35, 36]). The final property follows from the second characterisation in Theorem 4.11, equation (17) in the appendix, the equation of motion and the Propagation of Singularities Theorem for the wave front set, which in this case follows from the propagation of the polarisation set [12]. $\square$

One can also prove that the state spaces are locally physically equivalent [16] and that all quasi-free Hadamard states are locally quasi-equivalent [9]. Whether the latter remains true for all Hadamard states appears to be unknown.

We conclude this section with the remark that the functor $\mathbf{S} : \mathfrak{S}\mathfrak{P}\mathfrak{a}\mathfrak{c} \rightarrow \mathfrak{T}\mathfrak{M}\mathfrak{e}\mathfrak{c}$ defined by $SM \mapsto \mathcal{S}_{SM}$ and $\chi \mapsto \alpha_\chi^*$ (restricted to the relevant state space) is a locally covariant state space for the theory $\mathbf{B}$ [4].

4.3 The relative Cauchy evolution of the Dirac field and the stress-energy-momentum-tensor

Now that we have a locally covariant free Dirac field at our disposal we will investigate the idea of relative Cauchy evolution for this field and prove that it yields commutators with the stress-energy-momentum tensor. This result is completely analogous to the result for the free scalar field of [4].

Suppose that we have two objects $M_0 = (\mathcal{M}, g_0, SM_0, p_0)$ and $M_g = (\mathcal{M}, g, SM_g, p_g)$ in $\mathfrak{S}\mathfrak{P}\mathfrak{a}\mathfrak{c}$, where $\mathcal{M}$ is the same in both cases and such that outside a compact set $K \subset \mathcal{M}$ we have $g = g_0$, $SM_g = SM_0$ and $p_g = p_0$. Now let $N^\pm \subset M_0$ be causally convex open regions, each containing a Cauchy surface for M_0 , such that K lies to the future of N^- (i.e. $K \subset J^+(N^-) \setminus N^-$ in M_0 and hence also in M_g) and to the past of N^+ . We view $N^\pm$ as objects in $\mathfrak{S}\mathfrak{P}\mathfrak{a}\mathfrak{c}$ and consider the canonical morphisms $\iota_0^\pm : N^\pm \rightarrow M_0$ and $\iota_g^\pm : N^\pm \rightarrow M_g$. By the time-slice axiom, Proposition 4.8, these give rise to $*$ -isomorphisms $\beta_0^\pm : \mathcal{B}_{N^\pm} \rightarrow \mathcal{B}_{M_0}$ and $\beta_g^\pm : \mathcal{B}_{N^\pm} \rightarrow \mathcal{B}_{M_g}$. We then define

$$\beta_g := \beta_0^+ \circ (\beta_g^+)^{-1} \circ \beta_g^- \circ (\beta_0^-)^{-1}.$$

The $*$ -isomorphism $\beta_g : \mathcal{B}_{M_0} \rightarrow \mathcal{B}_{M_g}$ measures the change in an operator $A \in \mathcal{B}_{M_0}$ as it evolves to N^+ in the metric g instead of g_0 .²⁰ β_g can be extended to a $*$ -isomorphism of the algebra $\mathcal{F}_{M_0}$, where we fix the signs for the isomorphisms between the spinor bundles involved by identifying the double spinor bundles over $N^\pm \subset M_0$ and $N^\pm \subset M_g$. It represents the *relative Cauchy evolution* of the free Dirac field.

²⁰In [4] it seems the authors have the scattering of a state in mind as it passes through the perturbed metric, which leads them to consider the $*$ -isomorphisms $\beta_{g^{-1}}$ rather than β_g . When we take the variation w.r.t. g this gives rise to a sign.

We will want to compute the variation of the $*$ -isomorphism β_g as well as that of the action for the free Dirac field with respect to the metric g . For this purpose we will suppose that the compact set $K \subset \mathcal{M}$ has a contractible neighbourhood O which does not intersect either $N^\pm$. Let $\epsilon \mapsto g_\epsilon$ be a smooth curve from $[0, 1]$ into the space of Lorentzian metrics on $\mathcal{M}$ starting at g_0 and such that $g_\epsilon = g_0$ outside K for every ϵ . The spin bundle SM_ϵ must be trivial over the contractible region O . If we assume it to be diffeomorphic to SM_0 outside K we can simply take $SM_\epsilon = SM_0$ as a manifold and, choosing a fixed representation and matrices A, C , we obtain $DM_\epsilon = DM$.

The deformation of the spin structure is contained entirely in the spin frame projection $\pi_\epsilon : SM_0 \rightarrow FM_\epsilon$. Let E be a section of SM_0 over O and set $(e_\epsilon)_a := \pi_\epsilon(E)$. We require that e_ϵ varies smoothly with ϵ and that $(e_\epsilon)_a = (e_0)_a$ outside K . To show that projections π_ϵ with these properties exist we can apply the Gram-Schmidt orthonormalisation procedure to $(e_0)_a$ for all ϵ simultaneously. The assignment $E \mapsto e_\epsilon$ determines π_ϵ completely, using the intertwining properties. The family of frames e_ϵ determines principal fiber bundle isomorphisms $FM_\epsilon \rightarrow FM_0$ between the frame bundles by

$$\lambda_\epsilon : \{(e_\epsilon)_a\} \mapsto \{(e_0)_a\}$$

on K and extending it by the identity on the rest of $\mathcal{M}$. By definition f_ϵ intertwines the action of $\mathcal{L}_+^\dagger$ on the orthonormal frame bundles.

Remark 4.13 *There may be many deformations of the spin structure, i.e. many families of projections π_ϵ which satisfy our requirements. However, the variation of terms like $\langle v, P_\epsilon u \rangle$ will not depend on this choice. Indeed, if π'_ϵ is a different deformation of the spin structure, then $e'_\epsilon := \pi'_\epsilon(E) = R_{\Lambda_\epsilon} e_\epsilon = \pi_\epsilon(R_{S_\epsilon} E)$ for some smooth curve S_ϵ in $Spin_{1,3}^0$. However, using the invariance of $\langle, \rangle$ under the action of the gauge group $Spin_{1,3}^0$, the variation will be equal in both cases. (Also $\delta u = 0$ for every spinor u , because $D_\epsilon M = DM$.) In this sense the variation will only depend on the variation of the metric.*

4.3.1 The stress-energy-momentum tensor

The classical stress-energy-momentum tensor for the Dirac field is defined as a variation of the action $\mathcal{S} = \int_{\mathcal{M}} \mathcal{L}_D$, with the Lagrangian density (7), with respect to $g^{\mu\nu}(x)$:

$$T_{\mu\nu}(x) := \frac{2}{\sqrt{-\det g(x)}} \frac{\delta \mathcal{S}}{\delta g^{\mu\nu}(x)}, \quad (12)$$

where ψ is a free classical Dirac spinor, ψ^+ its adjoint. An explicit computation yields²¹

$$T_{\mu\nu} = \frac{i}{2} (\langle \psi^+, \gamma_{(\mu} \nabla_{\nu)} \psi \rangle - \langle \nabla_{(\mu} \psi^+, \gamma_{\nu)} \psi \rangle),$$

Here the brackets around indices denote symmetrisation as an idempotent operation and in the following indices between $|\dots|$ are to be excluded from the symmetrisation.

Following [17] we quantise the stress-energy-momentum tensor via a point-split procedure, i.e. we want to find a bi-distribution of scalar test-functions which reduces to $T_{\mu\nu}$ on the diagonal and which can be quantised in a straight-forward way. For this purpose we use a local spin frame E_A and recall that the components $\gamma_a^A{}_B$ of γ_a are constant. We define:

$$T_{ab}^s(x, y) := \frac{i}{2} \left(\langle \psi^+, E_A \rangle(x) \gamma_{(a|B|}^A \langle E^B, e_b^\mu \nabla_\mu \psi \rangle(y) - \langle e_a^\mu \nabla_{|\mu} \psi^+, E_A \rangle(x) \gamma_{b|}^A \langle E^B, \psi \rangle(y) \right),$$

reduces to $T_{ab} := e_a^\mu e_b^\nu T_{\mu\nu}$ in the limit $y \rightarrow x$. Performing a partial integration, $\int \nabla_\mu (e_a^\mu \langle v, u \rangle) = 0$, we can write T_{ab}^s as a bidistribution of scalar test-functions h_1, h_2 ,

$$\begin{aligned} T_{ab}^s(h_1, h_2) &= \frac{i}{2} \left(-\psi^+(E_A h_1) \gamma_{(a|B|}^A \psi(\nabla_{|\mu} (E^B e_b^\mu h_2)) \right. \\ &\quad \left. + \psi^+(\nabla_\mu (e_a^\mu E_{|A|} h_1)) \gamma_{b|}^A \psi(E^B h_2) \right). \end{aligned} \quad (13)$$

²¹For explicit computations we refer to section 4 of [18], who use a Lagrangian that differs from ours by a total derivative. Varying with respect to $g_{\mu\nu}$ would yield the opposite sign.

Equation (13) can be promoted to the quantised case by replacing ψ and ψ^+ by the components ψ_{SM} and ψ_{SM}^+ of the corresponding locally covariant quantum field. The expression (13) can be viewed as a formal expression for the same distribution with quantised field operators.

Proposition 4.14 *For all $f \in C_0^\infty(DM \oplus D^*M)$ and $h \in C_0^\infty(M)$ we have:*

$$\int_M [T_{ab}^s(x, x), B_{SM}(f)] h(x) d\text{vol}_g(x) = \frac{1}{2} \{ (\nabla_{(a} B_{SM})(\gamma_{b)}(SRf)h) - B_{SM}(\gamma_{(b} \nabla_{a)}(SRf)h) \},$$

where $\nabla_a := e_a^\mu \nabla_\mu$.

Proof. For $f = u \oplus v$ we use Proposition 4.6 to obtain:

$$\begin{aligned} \{B_{SM}(f), \psi_{SM}^+(E_A h)\} &= -i \int_M \langle v, S E_A h \rangle I = i \int_M \langle S_c v, E_A \rangle h I \\ \{B_{SM}(f), \psi_{SM}(\nabla_\mu E^B e_b^\mu h)\} &= -i \int_M \langle \nabla_\mu E^B e_b^\mu h, S u \rangle I = i \int_M \langle E^B, e_b^\mu \nabla_\mu S u \rangle h I \\ \{B_{SM}(f), \psi_{SM}^+(\nabla_\mu e_a^\mu E_A h)\} &= -i \int_M \langle v, S \nabla_\mu e_a^\mu E_A h \rangle I = -i \int_M \langle e_a^\mu \nabla_\mu S_c v, E_A \rangle h I \\ \{B_{SM}(f), \psi(E^B h)\} &= -i \langle E^B, S u \rangle h I. \end{aligned}$$

With equation (13), the commutation relations and $[AB, C] = A\{B, C\} - \{A, C\}B$ this implies

$$\begin{aligned} [T_{ab}^s(x, y), B_{SM}(f)] &= \frac{1}{2} \left\{ \psi_{SM}^+(E_A(x)) \gamma_{(a}^A{}_{|B|} \langle E^B, \nabla_{b)} S u \rangle(y) \right. \\ &\quad + \langle S_c v, E_A \rangle(x) \gamma_{(a}^A{}_{|B|} (\nabla_{b)} \psi_{SM})(E^B(y)) \\ &\quad - (\nabla_{(a} \psi_{SM}^+)(E_{|A|}(x)) \gamma_{b)}^A{}_{|B|} \langle E^B, S u \rangle(y) \\ &\quad \left. - \langle \nabla_{(a} S_c v, E_{|A|} \rangle(x) \gamma_{b)}^A{}_{|B|} \psi_{SM}(E^B(y)) \right\}. \end{aligned}$$

In this expression we are multiplying distributions with smooth functions, so we may take the coincidence limit yielding:

$$\begin{aligned} [T_{ab}^s(x, x), B_{SM}(f)] &= \frac{1}{2} \{ \psi_{SM}^+(\gamma_{(a} \nabla_{b)}(S u)(x)) + \nabla_{(b} \psi_{SM}(S_c v \gamma_{a)}(x)) \\ &\quad - \nabla_{(a} \psi_{SM}^+(\gamma_{b)} S u(x)) - \psi_{SM}(\nabla_{(a} (S_c v) \gamma_{b)}(x)) \} \\ &= \frac{-1}{2} \{ \nabla_{(a} B_{SM}(\gamma_{b)} S R f(x)) - B_{SM}(\gamma_{(b} \nabla_{a)}(S R f)(x)) \}, \end{aligned}$$

from which the result follows. $\square$

This result can be written for spinors and cospinors separately as:

$$\begin{aligned} \int_M [T_{ab}^s(x, x), \psi_{SM}(v)] h(x) d\text{vol}_g(x) &= \frac{1}{2} \{ \nabla_{(a} \psi_{SM}((S_c v) \gamma_{b)} h) - \psi_{SM}(\nabla_{(a} (S_c v) \gamma_{b)} h) \} \\ \int_M [T_{ab}^s(x, x), \psi_{SM}^+(u)] h(x) d\text{vol}_g(x) &= \frac{-1}{2} \{ \nabla_{(a} \psi_{SM}^+(\gamma_{b)} S u h) - \psi_{SM}^+(\gamma_{(a} \nabla_{b)}(S u) h) \}. \end{aligned}$$

4.3.2 Relative Cauchy evolution

To compute the relative Cauchy evolution explicitly we first note that the isomorphism β_g can be characterised in terms of its action on the generators $B_{M_0}(f)$ of $\mathcal{F}_{M_0}$ as follows:

Proposition 4.15 *For $f \in C_0^\infty(DN^+ \oplus D^*N^+)$ we have $\beta_g B_0(f) = B_0(T_g f)$, where*

$$T_g f = P_g \phi_+ S_g P_0 \phi_- S_0 f.$$

Here the subscripts on B , P and S indicate whether they are the objects defined on M_0 or M_g and the smooth functions $\phi_\pm$ are such that $\phi_\pm \equiv 1$ to the past of some Cauchy surface in $N^\pm$ and $\phi_\pm \equiv 0$ to the future of some other Cauchy surface in $N^\pm$.

Proof. Note that $\beta_g^- \circ (\beta_0^-)^{-1} B_0(\tilde{f}) = B_g(\tilde{f})$ for any $\tilde{f} \in C_0^\infty(DN^- \oplus D^*N^-)$. Similarly, for $f' \in C_0^\infty(DN^+ \oplus D^*N^+)$ we have $\beta_0^+ \circ (\beta_g^+)^{-1} B_g(f') = B_0(f')$. The functions $\phi_\pm, 1 - \phi_\pm$ have been chosen appropriately in order to apply equation (11) in Proposition 4.8. We then have $B_0(\tilde{f}) = B_0(f)$, where $\tilde{f} := -P_0\phi_-S_0f$. Notice that $\tilde{f}$ indeed has a compact support in N^- . Similarly, $B_g(f) = B_g(f')$, where $f' := -P_g\phi_+S_g\tilde{f}$ has support in N^+ . Hence, for $f' = T_gf$: $\beta_g B_0(f) = \beta_g B_0(\tilde{f}) = \beta_0^+ \circ (\beta_g^+)^{-1} B_g(\tilde{f}) = \beta_0^+ \circ (\beta_g^+)^{-1} B_g(f') = B_0(f')$. $\square$

On each spin spacetime $M_\epsilon = (\mathcal{M}, g_\epsilon, SM_0, \pi_\epsilon)$ we can now quantise the Dirac field and obtain relative Cauchy evolutions $\beta_\epsilon := \beta_{g_\epsilon}$ on $\mathcal{F}_{N^+}$ as before.

Proposition 4.16 *Writing $\delta := \partial_\epsilon|_{\epsilon=0}$ we have for all $f \in C_0^\infty(DN^+ \oplus D^*N^+)$:*

$$\delta(\beta_\epsilon B_0(f)) = B_0(R(\delta \nabla_\epsilon) S_0 f).$$

Proof. Using the fact that B_0 is a C^* -algebra-valued distribution and Proposition 4.15 we find:

$$\begin{aligned} \delta(\beta_\epsilon B_0(f)) &= \delta(B_0(P_\epsilon \phi_+ S_\epsilon P_0 \phi_- S_0 f)) = B_0(\delta(P_\epsilon \phi_+ S_\epsilon) P_0 \phi_- S_0 f) \\ &= B_0(\delta(P_\epsilon) \phi_+ S_0 P_0 \phi_- S_0 f) + B_0(P_0 \phi_+ \delta(S_\epsilon) P_0 \phi_- S_0 f). \end{aligned}$$

Now, because $P_0 \phi_- S_0 f \in C_0^\infty(DN^- \oplus D^*N^-)$ we see that $\delta(S_\epsilon) P_0 \phi_- S_0 f$ vanishes on $J^-(N^-)$ and that $\phi_+ \delta(S_\epsilon) P_0 \phi_- S_0 f$ has compact support. Because B_0 solves the Dirac equation we conclude that the second term vanishes. The first term can be rewritten using equation (11), which yields $S_0 f = -S_0 P_0(\phi_- S_0 f)$ and hence:

$$\delta(\beta_\epsilon B_0(f)) = -B_0(\delta(P_\epsilon) \phi_+ S_0 f) = -B_0(\delta(P_\epsilon) S_0 f).$$

For the last equality we used the fact that $\delta(P_\epsilon)$ is supported in K , where $\phi_+ \equiv 1$. Recall that $P = (-i\nabla + m) \oplus (i\nabla + m)$ to get the final result. $\square$

To compute the variation of the Dirac operator we may work in a local frame on O , where it is supported. Because the Dirac adjoint map is independent of ϵ we only need to compute this variation either for spinors or for cospinors:

Lemma 4.17 *For $v \in C_0^\infty(D^*M)$ we have $\delta(\nabla)v = (\delta(\nabla)v)^+$.*

Proof. Because the adjoint operation is continuous we have:

$$\delta(\nabla)v = \partial_\epsilon \nabla_\epsilon v|_{\epsilon=0} = \partial_\epsilon (\nabla_\epsilon v)^+|_{\epsilon=0} = (\partial_\epsilon \nabla_\epsilon v^+|_{\epsilon=0})^+ = (\delta(\nabla)v^+)^+.$$

$\square$

It is interesting to note that only the variation of the Dirac operator is of importance for the variation of the relative Cauchy evolution, just like for the stress-energy-momentum tensor (cf. [18]). It will also turn out that the variation only depends on the variation of the metric and not on the other freedom in the variation of the orthonormal frame, even though we are now acting on it with the C^* -algebra-valued field (cf. Remark 4.13). This will follow from the proof of the following theorem, for which we refer to appendix B.

Theorem 4.18 *For a double test-spinor $f \in C_0^\infty(DM_0 \oplus D^*M_0)$ and $x \in K$:*

$$\frac{\delta}{\delta g^{\alpha\beta}(x)} (\beta_g B_0(f)) = -B_0 \left(\frac{\delta}{\delta g^{\alpha\beta}(x)} P_g S_0 f \right) = \frac{-i}{2} e_\alpha^a e_\beta^b [T_{ab}^s(x, x), B_0(f)]. \quad (14)$$

This result compares well with the scalar field case, Theorem 4.3 in [4].²² As particular cases we obtain for ψ and ψ^+ :

$$\begin{aligned} \frac{\delta}{\delta g^{\alpha\beta}(x)} (\beta_g \psi(v)) &= \frac{-i}{2} e_\alpha^a e_\beta^b [T_{ab}^s(x, x), \psi(v)], \\ \frac{\delta}{\delta g^{\alpha\beta}(x)} (\beta_g \psi^+(u)) &= \frac{-i}{2} e_\alpha^a e_\beta^b [T_{ab}^s(x, x), \psi^+(u)]. \end{aligned}$$

It follows that the same result also holds for products and sums of smeared field operators.

²²The sign explained in the footnote on page 23 cancels the sign due to the variation w.r.t. $g^{\alpha\beta}$ instead of $g_{\alpha\beta}$.

5 Conclusions

A rigorous formulation of quantum field theories in curved spacetime, going beyond the well-known scalar field, is a prerequisite for constructing more realistic cosmological models as well as for improving our understanding of quantum field theory in Minkowski spacetime. The main purpose of this paper was to present the free Dirac field in a four dimensional globally hyperbolic spacetime as a locally covariant quantum field theory in the sense of [4] and to compute the relative Cauchy evolution of this field, obtaining commutators with the stress-energy-momentum tensor in analogy with the free real scalar field. We achieved this in a representation independent way and in a functorial, and therefore manifestly covariant, framework.

We established some basic properties of the locally covariant free Dirac field and remarked on the quantisation of Majorana spinors. We also provided a detailed discussion of Hadamard states, closing any gaps in the existing proofs of the equivalence of the definitions in terms of the series expansion of their two-point distribution and a microlocal condition, respectively.

Furthermore we argued that the observable part of the theory is uniquely determined by the relations between adjoints, charge conjugation and the Dirac operator, although the geometric constructions themselves may not be unique due to the cohomological properties of the category of spin spacetime. On a mathematical level we have consistently replaced a single spin spacetime SM by the category $\mathfrak{Spac}$ of such spacetimes, and the differential geometry on SM by the corresponding functorial descriptions. On a physical level, however, we should not conclude from this that $\mathfrak{Spac}$ is now the physical arena in which our system lives, instead of a collection of systems. (See Ch.1 of [36] for more detailed philosophical remarks on the interpretation of the locally covariant approach.)

Acknowledgements

I would like to thank Chris Fewster for suggesting to use the cohomological language in subsection 3.4 and for bringing the problem of computing the relative Cauchy evolution for the Dirac field to my attention. I would also like to thank Romeo Brunetti for correcting some of my misconceptions in the early stages of this computation. Much of this work was performed as part of my PhD-thesis at the University of York and I would also like to thank the University of Trento for their kind hospitality during my visit in October 2007. Furthermore this research was supported by the German Research Foundation (Deutsche Forschungsgemeinschaft (DFG)) through the Institutional Strategy of the University of Göttingen and the Graduiertenkolleg 1493 "Mathematische Strukturen in der modernen Quantenphysik".

A Results on microlocal analysis

In this appendix we will list some results concerning the microlocal analysis of distributions. For a detailed treatment of scalar distributions we refer to [24], whereas Hilbert and Banach-space-valued distributions are treated in [39, 36]. More details concerning distributional sections of vector bundles can be found in e.g. [2, 34, 12, 36].

Before we discuss distributional sections of vector bundles we first consider the scaling limit of a distribution in an open set of $\mathbb{R}^n$:

Definition A.1 *Let O be a convex open region $O \subset \mathbb{R}^n$ containing 0. For all $\lambda > 0$ we define the scaling map $\delta_\lambda: O \rightarrow O$ by $\delta_\lambda(x) := \lambda x$.*

Let u be a distribution on a convex open region $O \subset \mathbb{R}^n$ containing 0. The scaling degree d of u at 0 is defined as $d := \inf \{ \beta \in [-\infty, \infty) \mid \lim_{\lambda \rightarrow 0} \lambda^\beta \delta_\lambda^ u = 0 \}$, where $(\delta_\lambda^* u)(f) := \lambda^{-n} u(f \circ \delta_\lambda^{-1})$.*

If $u^0 := \lim_{\lambda \rightarrow 0} \lambda^d \delta_\lambda^ u$ exists we call it the scaling limit of u at 0.*

Note that the scaling limit may fail to exist (e.g. $u(x) = \log|x|$) or it may vanish (e.g. if $0 \notin \text{supp}(u)$). On a manifold, we will only consider scaling limits in a certain choice of local coordinates. How this limit depends on this choice of coordinates will not be relevant for us.

We now prove the following result:²³

Proposition A.2 *Let u be a distribution on a convex open region $O \subset \mathbb{R}^n$ containing 0 with scaling limit u^0 at 0. Then*

$$\{0\} \times \pi_2(WF(u^0)) \subset WF(u),$$

where π_2 denotes the projection on the second coordinate.

Proof. Suppose that $(0, \xi_0) \notin WF(u)$ with $\xi_0 \neq 0$. We will prove that $(x, \xi_0) \notin WF(u^0)$ for all x . By assumption we can choose $\chi \in C_0^\infty(O)$ and an open conic neighbourhood $\Gamma \subset \mathbb{R}^n$ of ξ_0 such that $\chi \equiv 1$ on a neighbourhood of 0 and $\text{supp}(\chi) \times \Gamma \cap WF(u) = \emptyset$. We set $v := \chi u$ and $v^\lambda := \lambda^d \delta_\lambda^* v$, where d is the scaling degree of u at 0. Notice that $WF(v) \cap T_0^* O = WF(u) \cap T_0^* O$ and $u^0 := \lim_{\lambda \rightarrow 0} v^\lambda$, so without loss of generality we may prove the result with v replacing u and we can view the v^λ as compactly supported distributions on all of $\mathbb{R}^n$.

Notice that for $\lambda > 0$ we have $\delta_\lambda^* u^0 = \lambda^{-d} u^0$, i.e. u^0 is a homogeneous distribution and therefore it is tempered ([24] Theorem 7.1.18). We now prove that v^λ converges to u^0 in the sense of tempered distributions on $\mathbb{R}^n$. For this we first write $v = \sum_{|\alpha| \leq r} (-1)^{|\alpha|} \partial^\alpha v_\alpha$, where r is the order of v and the v_α are compactly supported distributions of order 0 (see [24] Section 2.1). Note that $\sum_{|\alpha| < d-n} (-1)^{|\alpha|} \partial^\alpha v_\alpha$ converges to 0 in $\mathcal{S}$, because for every $|\alpha| < d-n$ and $\phi \in \mathcal{S}(\mathbb{R}^n)$ we have

$$|((-1)^{|\alpha|} \partial^\alpha v_\alpha)^\lambda(\phi)| = \lambda^{d-n} |v_\alpha(\partial^\alpha(\phi \circ \delta_\lambda^{-1}))| \leq \lambda^{d-n-|\alpha|} C \sup |\partial^\alpha \phi|$$

which converges to 0 as $\lambda \rightarrow 0$. We then set $w := \sum_{d-n \leq |\alpha| \leq r} (-1)^{|\alpha|} \partial^\alpha v_\alpha$, so that $\lim_{\lambda \rightarrow 0} w^\lambda = u^0$ as distributions. By the Uniform Boundedness Principle this implies

$$|w^\lambda(\phi)| \leq C \sum_{|\alpha| \leq r} \sup |\partial^\alpha \phi|, \quad \text{supp}(\phi) \subset B_1, \quad (15)$$

for some $C, r > 0$, where B_1 is the (Euclidean) unit ball and $0 < \lambda \leq 1$. In fact, for $\lambda \geq 1$ we also have

$$|w^\lambda(\phi)| = \lambda^{d-n} |w(\phi \circ \delta_\lambda^{-1})| \leq C \sum_{d-n \leq |\alpha| \leq r} \lambda^{d-n-|\alpha|} \sup |\partial^\alpha \phi| \leq C \sum_{d-n \leq |\alpha| \leq r} \sup |\partial^\alpha \phi|,$$

so the estimate (15) holds for all $\lambda > 0$.

Now let $\phi \in \mathcal{S}(\mathbb{R}^n)$ be a function of rapid decrease and choose a partition of unity on $\mathbb{R}^n$ as follows. We let $\chi_0 \in C_0^\infty(\mathbb{R}^n)$ be positive such that $\chi \equiv 1$ on B_1 and $\chi(x) = 0$ when $\|x\| \geq 2$. We then set $\chi_m(x) := \chi_0(2^{-m}x) - \chi_0(2^{1-m}x)$ and note that:

$$\text{supp}(\chi_{m \geq 1}) \subset \{x \mid 2^{m-1} \leq \|x\| \leq 2^{m+1}\}, \quad \sum_{m=0}^{\infty} \chi_m = 1,$$

where the sum is finite near every point. We define $\phi_m := \chi_m \phi$ and $\mu_m := 2^{-m-1}$ and rescale ϕ_m in order to apply the estimate (15):

$$\begin{aligned} |w^\lambda(\phi_m)| &= \mu_m^{d-n} \left| w^{\lambda/\mu_m} \left(\phi_m \left(\frac{\cdot}{\mu_m} \right) \right) \right| \leq C \sum_{|\alpha| \leq r} \mu_m^{d-n-|\alpha|} \sup \left| (\partial^\alpha \phi_m) \left(\frac{\cdot}{\mu_m} \right) \right| \\ &\leq C_1 \sum_{|\alpha| \leq r} \sum_{|\beta| \leq r+n-d} \sup_{\mathbb{R}^n} |x^\beta \partial^\alpha \phi_m|, \quad m \geq 0, \end{aligned} \quad (16)$$

²³A similar result was also claimed in [34], but we find the proof unconvincing. In particular, when localising the scaling limit with a test-function χ_0 and estimating

$$\widehat{\chi_0 u^0}(\xi) = \lim_{\lambda \rightarrow 0} \lambda^{d-n} u \left(\chi_0 \left(\frac{\cdot}{\lambda} \right) e^{-i \frac{\xi}{\lambda} \cdot} \right)$$

the test-function $\chi_0(\frac{\cdot}{\lambda})$ becomes singular in the limit $\lambda \rightarrow 0$. The quoted reference pays insufficient attention to this issue.

where the last line uses $\mu_m^{d-n-|\alpha|} \leq (4\|x\|)^{|\alpha|+n-d}$ for $m \geq 1$, which follows from $d-n \leq |\alpha|$ and the support properties of χ_m . (For $m = 0$ we simply estimate $\mu_0^{d-n-|\alpha|}$ by a constant to arrive at the last line of (16).) We now note that $\max_\alpha \sup_x |\partial^\alpha \chi_m| \leq c$ for some c independent of m , as the derivatives only bring out extra factors of $2^{-m} \leq 1$. Moreover, for $m \geq 0$ we notice that $\chi_{m+1} + \chi_m + \chi_{m-1} \equiv 1$ on $\text{supp}(\chi_m)$, where we define $\chi_{-1} := 0$. Therefore (16) leads to

$$|w^\lambda(\phi_m)| \leq C_2 \sum_{|\alpha| \leq r} \sum_{|\beta| \leq r+n-d} \sup_{\mathbb{R}^n} |x^\beta \partial^\alpha \phi| (\chi_{m+1} + \chi_m + \chi_{m-1})$$

and summing over $m \geq 0$ then gives:

$$|w^\lambda(\phi)| \leq 3C_2 \sum_{|\alpha| \leq r} \sum_{|\beta| \leq r+n-d} \sup_{\mathbb{R}^n} |x^\beta \partial^\alpha \phi|.$$

This shows that $w^\lambda(\phi)$ can be estimated by a seminorm on $\mathcal{S}(\mathbb{R}^n)$ uniformly in λ . It then follows that $w^\lambda \rightarrow u^0$ and hence $v^\lambda \rightarrow u^0$ as tempered distributions. Indeed, for any $\phi \in \mathcal{S}(\mathbb{R}^n)$ and $\epsilon > 0$ we can choose $\phi' \in C_0^\infty(\mathbb{R}^n)$ and $\lambda_0 > 0$ such that $|w^\lambda(\phi - \phi')| < \frac{\epsilon}{2}$ for all $\lambda > 0$ and $|w^\lambda(\phi')| < \frac{\epsilon}{2}$ for all $\lambda < \lambda_0$.

Fourier transformation is a continuous operation on tempered distributions, so we can compute:

$$|\widehat{u^0}(\xi)| = \lim_{\lambda \rightarrow 0} \lambda^{d-n} \left| \widehat{v} \left(\frac{\xi}{\lambda} \right) \right| \leq C_N \lim_{\lambda \rightarrow 0} \lambda^{d-n} \left\| \frac{\xi}{\lambda} \right\|^{-N} = C_N \|\xi\|^{-N} \lim_{\lambda \rightarrow 0} \lambda^{N+d-n}$$

for all ξ in Γ , all $N \in \mathbb{N}$ and suitable $C_N > 0$. For $N > n - d$ the limit yields $\widehat{u^0}(\xi) = 0$ near ξ_0 . We then apply Theorem 8.1.8 in [24], which says that for a homogeneous distribution we have for all $x \neq 0$ that $(x, \xi_0) \in WF(u^0)$ if and only if $(\xi_0, -x) \in WF(\widehat{u^0})$ and also $(0, \xi_0) \in WF(u^0)$ if and only if $\xi_0 \in \text{supp}(\widehat{u^0})$. $\square$

For a distribution u with values in a Banach space $\mathcal{B}$ one can define the wave front set by using estimates of the norm $\|u(\chi e^{i\xi \cdot})\|$, which replace the corresponding estimates of the absolute value $|u(\chi e^{i\xi \cdot})|$ for scalar distributions [39]. Alternatively, one can use the following equivalent characterisation ([36], Theorem A.1.4):

$$WF(u) = \overline{\cup_{l \in \mathcal{B}'} WF(l \circ u)} \setminus \mathcal{Z}. \quad (17)$$

A similar idea works for a distributional section u of a vector bundle $\mathcal{V} = O \times \mathbb{R}^m$ over a contractible region O of $\mathbb{R}^n$. Indeed, using a basis e_i for $\mathbb{R}^m$ with dual basis e^i we can identify u with a distribution $\tilde{u}$ on O with values in $\mathcal{B} \otimes (\mathbb{R}^m)'$, where the correspondence is given by

$$\tilde{u}(h) := \sum_{i=1}^m u(h e_i) \otimes e^i, \quad u \left(\sum_{i=1}^m f^i e_i \right) = \sum_{i=1}^m \langle \tilde{u}(f^i), e_i \rangle,$$

where $\langle \cdot, \cdot \rangle$ denotes the canonical pairing of $\mathbb{R}^m$ with the second factor of $\mathcal{B} \otimes (\mathbb{R}^m)'$. We set by definition $WF(u) := WF(\tilde{u})$.

Equation (17) allows a straightforward generalisation of many results for scalar distributions on open sets of $\mathbb{R}^n$ to Banach-space-valued distributional sections of a vector bundle over regions over $\mathbb{R}^n$. Moreover, by showing how these results transform under changes of coordinates they can be formulated for vector bundles on a manifold. We list a number of these results in the following Theorem (cf. [24, 36]):

Theorem A.3 *If u, v are distributional sections of a complex vector bundle $\mathcal{V}$ over the spacetime M with values in the Banach space $\mathcal{B}$, then:*

1. *sing supp(u) is the projection of $WF(u)$ on the first variable,*
2. *$u \in C^\infty(\mathcal{V}, \mathcal{B})$ if and only if $WF(u) = \emptyset$,*

3. $WF(u + v) \subset WF(u) + WF(v)$,
4. if P is a linear partial differential operator on $\mathcal{V}$ with smooth coefficients and (matrix-valued) principal symbol²⁴ $p(x; \xi)$, then $WF(Pu) \subset WF(u) \cup WF(Pu) \cap \Omega_P$, where $\Omega_P := \{(x; \xi) \in T^*M \mid \xi \neq 0, \det p(x; \xi) = 0\}$,
5. if $x \in M$, $\phi: U \rightarrow \mathbb{R}^n$ is a local trivialisation on a convex neighbourhood U with $\phi(x) = 0$ and $(\phi^{-1})^*u$ has a scaling limit u^0 at 0, then $\phi^*({0} \times \pi_2(WF(u^0))) \subset WF(u) \cap T_x^*M$.

In the last item, the scaling limit depends not just on the choice of coordinates, but also on the choice of a frame e_i of $\mathcal{V}$ over U and we let the scaling maps δ_λ act on sections of $\mathcal{V}$ componentwise: $(\sum_i f^i e_i) \circ \delta_\lambda^{-1} = \sum_i (f^i \circ \delta_\lambda^{-1}) e_i$.

In the particular case where $\mathcal{B}$ is a Hilbert space we also have (see [39, 36]):

Theorem A.4 *Let $\mathcal{H}$ be a Hilbert space and $\mathcal{V}_i$, $i = 1, 2$, two finite dimensional (complex) vector bundles over smooth n_i dimensional spacetimes M_i with complex conjugations J_i , i.e. the J_i are antilinear, base-point preserving bundle isomorphisms $J_i: \mathcal{V}_i \rightarrow \mathcal{V}_i$ such that $J_i^2 = -\text{id}$. Let u_i , $i = 1, 2$, be two $\mathcal{H}$ -valued distributional sections of $\mathcal{V}_i$ and let w_{ij} be the distributional sections of the vector bundle $\mathcal{X}_i \boxtimes \mathcal{X}_j$ over $M_i \times M_j$ determined by $w_{ij}(f_1 \boxtimes f_2) := \langle u_i(J_i f_1), u_j(f_2) \rangle$. Then*

$$(x, \xi) \in WF(u_1) \Leftrightarrow (x, -\xi; x, \xi) \in WF(w_{11})$$

and

$$WF(w_{ij}) \subset -(WF(u_i) \cup \mathcal{Z}) \times (WF(u_j) \cup \mathcal{Z}),$$

where $\mathcal{Z}$ denotes the zero-section.

Finally we establish some results on the wave front sets of advanced and retarded fundamental solutions $E^\pm$ (for their existence and uniqueness we refer to [2]) and $S^\pm, S_c^\pm$. These results are analogous to Theorem 6.5.3 of [15], but now for operators in a vector bundle. Note that for distributional sections of vector bundles there is a Propagation of Singularities Theorem, which follows from the propagation of the polarisation set [12].

Theorem A.5 *Let $E^\pm$ be the advanced $(-)$ and retarded $(+)$ fundamental solutions for a normally hyperbolic operator P acting on the sections of a vector bundle DM over a globally hyperbolic spacetime $M = (\mathcal{M}, g)$ of dimension $n \geq 2$. Then*

$$\begin{aligned} WF(E^\pm) &= \{(x, \xi; y, \eta) \in T^*M^{\times 2} \setminus \mathcal{Z} \mid x \in J^\pm(y), x \neq y, (x, -\xi) \sim (y, \eta)\} \\ &\quad \cup \{(x, -\xi; x, \xi) \in T^*M^{\times 2} \setminus \mathcal{Z} \mid (x, \xi) \in T^*M \setminus \mathcal{Z}\} \\ &=: A^\pm \cup B \end{aligned} \tag{18}$$

where $\mathcal{Z}$ is the zero-section and $(x, \xi) \sim (y, \eta)$ if and only if there is a light-like geodesic γ from x to y to which ξ and η are cotangent such that they are each others parallel transport along γ .

Proof. The first part of this proof follows closely the proof of [32].

We start by reducing the problem to a local one as follows. The principal symbol of P is $p(x, \xi) = g_{\mu\nu}(x) \xi^\mu \xi^\nu I$, where I is the identity operator on DM , so by the Propagation of Singularities Theorem, the singularities of $E^\pm$ propagate along light-like geodesics by parallel transport. By definition the points in set $A^\pm$ are invariant under the same parallel transport. Now consider a point $p := (x, \xi; y, \eta)$ with $x \neq y$. If $\xi = \eta = 0$ then P is not contained in any set on either side of the equality, so we may assume $\xi \neq 0$ (the case $\eta \neq 0$ is analogous). Let S be a spacelike Cauchy surface through y and propagate (x, ξ) along the light-like geodesic γ towards S . If γ ends at S in $x' \neq y$ then P is not contained in $A^\pm$ or B , nor is it contained in $WF(E^\pm)$, because $E(x', y) = 0$ when x' and y are spacelike, so it cannot have any singularities there. If γ ends at y , on the other hand, we can find a point $p' := (x', \xi'; y, \eta)$, where x' on γ is in any given causally

²⁴See [2] for the definition of the principal symbol.

convex neighbourhood of y and ξ' is the parallel transport of ξ along γ to x' . Then $p' \in WF(E^\pm)$ if and only if $p \in WF(E^\pm)$ and $p' \in A^\pm$ if and only if $p \in A^\pm$. Hence, it suffices to prove the claim locally.

On a sufficiently small causally convex domain $O \subset \mathcal{M}$ we can find for every $k \in \mathbb{N}$ a C^k -section W^k of $DM \boxtimes D^*M$ on $O^{\times 2}$ such that ([2] Proposition 2.5.1):

$$E^\pm(x, y) = \sum_{j=0}^{k+1} V_j(x, y) f^*(1 \otimes R^\pm(2 + 2j, \cdot))(x, y) + W^k(x, y). \quad (19)$$

Here the Hadamard coefficients V_j are uniquely defined smooth sections of $DM \boxtimes D^*M$ on $O^{\times 2}$, $R^\pm(\alpha, y)$ are the retarded (+) and advanced (−) Riesz distributions (or rather distribution densities) on Minkowski spacetime and they are pulled back by the smooth diffeomorphism $f: O^{\times 2} \rightarrow TO$ defined by $(x, y) \mapsto (x, \exp_x^{-1}(y))$. This means we use Riemannian normal coordinates for y centered on x , which is well-defined because O is causally convex. The Riesz distributions have many useful properties, of which we will only use for all $j \geq 0$:

$$\begin{aligned} WF(R^\pm(2j + 2, \cdot)) &= \{(x, \xi) \in T^*M_0 \setminus \mathcal{Z} \mid x = 0 \text{ or } x^2 = 0, x \in J^\pm(0), \xi \parallel x\} \\ R^\pm(2 + 2j, \lambda x) &= \lambda^{2+2j-n} R^\pm(2 + 2j, x), \quad \lambda > 0 \end{aligned} \quad (20)$$

(These can be proved using [2] Proposition 1.2.4 items 4, and 5, $\square^{j+1} R^\pm(2 + 2j, \cdot) = \delta$ and the wave front sets of the distinguished parametrices as determined in [15].) Hence, for all $j \in \mathbb{N}$:

$$\begin{aligned} WF(f^*(1 \otimes R^\pm(2 + 2j, \cdot))) &= f^*(WF(1 \otimes R^\pm(2 + 2j, \cdot))) = f^*(\mathcal{Z}|_O \times WF(R^\pm(2 + 2j, \cdot))) \\ &= \{(x, \xi; y, \eta) \mid (\xi, \eta) = df^T(0, \eta') \text{ for some} \\ &\quad (\exp_x^{-1}(y), \eta') \in WF(R^\pm(2 + 2j, \cdot))\}, \\ &= (A^\pm \cup B) \cap T^*O^{\times 2}, \end{aligned} \quad (21)$$

where df^T is the transpose of the derivative df at (x, y) . The last equality uses the wave front set of the Riesz distributions in equation (20) and the properties of Riemannian normal coordinates (cf. [32]). It follows that $WF(E^\pm|_{O^{\times 2}}) \subset (A^\pm \cup B) \cap T^*O^{\times 2}$, because for each order of differentiation N we can choose a sufficiently high order k in equation (19) to make the required estimate in the definition of the wave front set.

We can prove the opposite inclusion, if we can show that the wave front set of the finite sum in (19) also contains $(A^\pm \cup B) \cap T^*O^{\times 2}$, which we will do using scaling limits (cf. [34]). First we may employ the Riemannian normal coordinates $f: O^{\times 2} \rightarrow TO$ as above. Next we may assume that O is also a contractible coordinate neighbourhood, so we can consider local coordinates $\phi: O \rightarrow \mathbb{R}^n$ on O and the associated coordinate map $d\phi$ on TO . Moreover, we can choose ϕ in such a way that $\phi(x_0) = 0$ for an arbitrarily given $x_0 \in O$. The composition $d\phi \circ f$ then defines coordinates on $O^{\times 2}$ such that $(x_0, x_0) \mapsto 0 \in \mathbb{R}^{2n}$. Using a frame E_A for $DM|_O$ and the dual frame E^B we can express the terms in the sum of equation (19) in the local coordinates $d\phi \circ f$ as $V_{jB}^A(x, y) R^\pm(2 + 2j, y)$. From equation (20) we then find the scaling behaviour

$$\delta_\lambda^*(V_{jB}^A(x, y) R^\pm(2 + 2j, y)) = \lambda^{2+2j-n} (V_{jB}^A(\lambda x, \lambda y) R^\pm(2 + 2j, y))$$

for all $\lambda > 0$. In the scaling limit only the lowest order term survives:

$$\lim_{\lambda \rightarrow 0} \lambda^{n-2} (\delta_\lambda \circ f^{-1} \circ d\phi^{-1})^* E(x, y) = V_{0B}^A(0, 0) R(2, y) E^B(x) E_A(y) = R(2, y) E^A(x) E_A(y),$$

where we wrote $R(2, y) := R^-(2, y) - R^+(2, y)$ and we used the explicit expression $V_{0B}^A(x, x) = \delta_B^A$ ([2] Lemmas 2.2.2 and 1.3.17).

Now, the last item of Theorem A.3 (which follows from Proposition A.2) implies that $WF(E) \supset (d\phi \circ f)^*(\{(0, 0)\} \times \pi_2(WF(1 \otimes R(2, \cdot))))$, because $E^A(x) E_A(y)$ is smooth and not identically vanishing. From equation (20) and the support properties of $R^\pm(2, \cdot)$ we easily compute $\pi_2(WF(1 \otimes$

$R(2, \cdot)) = \{(0, \xi) \mid \xi^2 = 0\}$. Pulling this back to $O^{\times 2}$ and using the properties of Riemannian normal coordinates yields

$$WF(E) \supset \{(x_0, -\xi; x_0, \xi) \mid \xi^2 = 0\}.$$

Because E is a bi-solution to the wave equation we can apply the Propagation of Singularities Theorem to find that $WF(E) \supset A^+ \cup A^-$ on $O^{\times 2}$ and from the support properties of E^+ and E^- we then conclude that $WF(E^\pm) \supset A^\pm$. Finally, $WF(E^\pm) \supset WF(PE^\pm) = WF(\delta) = B$. This completes the proof. $\square$

Corollary A.6 *In the notation of Theorem A.5, $WF(E) = \overline{A^+ \cup A^-} \setminus \mathcal{Z}$.*

Proof. By Theorem A.5 and the support properties of $E^\pm$ we have $WF(E) = A^+ \cup A^-$ away from the diagonal. The inclusion $\supset$ then follows from the closedness of the wave front set. For the opposite inclusion we consider a point on the diagonal and use the Propagation of Singularities Theorem to find an approximating sequence of points off the diagonal. $\square$

Proposition A.7 *For the fundamental solutions of the Dirac equation we have, in the notation of Theorem A.5: $WF(S^\pm) = WF(S_c^\pm) = A^\pm \cup B$ and $WF(S) = WF(S_c) = \overline{A^+ \cup A^-} \setminus \mathcal{Z}$.*

In other words, $WF(S^\pm) = WF(S_c^\pm) = WF(E^\pm)$ and $WF(S) = WF(S_c) = WF(E)$.

Proof. Because $S^\pm = (i\nabla + m)E^\pm$ and $S_c^\pm = (-i\nabla + m)E^\pm$ (see [14]) we immediately find $WF(S^\pm) \subset WF(E^\pm)$ and $WF(S_c^\pm) \subset WF(E^\pm)$. Similarly $WF(S) \subset WF(E)$ and $WF(S_c) \subset WF(E)$. Now suppose that $WF(S) = WF(S_c) = WF(E) = \overline{A^+ \cup A^-}$, which we will prove below. By the support properties of the fundamental solutions we then find that away from the diagonal $WF(S^\pm) = WF(S_c^\pm) = A^\pm$, whereas on the diagonal $WF(E^\pm) = B \supset WF(S^\pm) \supset WF(PS^\pm) = WF(\delta) = B$ and similarly for cospinors.

To complete the proof we need to show that $WF(S) \supset WF(E)$ and $WF(S_c) \supset WF(E)$, for which we adapt (and correct) an idea of [22]. We prove the case of S , because the other case follows by taking adjoints (cf. Theorem 3.10). Further note that it is sufficient to prove the claim on the diagonal, because the Propagation of Singularities Theorem applies both to E and to S . Now suppose that $(x, -\xi; x, \xi) \in WF(E) \setminus WF(S)$. We will derive a contradiction as follows. For every time-like, future pointing normalised vector $n_0 \in T_x M$ we can find a smooth spacelike Cauchy surface C through x such that n_0 is normal to C . We let n denote the future pointing normal vector field on C and $\iota: C \rightarrow M$ the canonical injection. By [14] Proposition 2.4c) we can restrict S to $C^{\times 2}$ to find $S|_{C^{\times 2}} = -i\delta\eta$ and in particular $(x, -d\iota_x^T(\xi); x, d\iota_x^T(\xi)) \in WF(S|_{C^{\times 2}})$. By (a component version of) [24] Theorem 8.2.4, on the other hand:

$$WF(S|_{C^{\times 2}}) \subset (\iota \times \iota)^*(WF(S)) = \{(x, d\iota_x^T(\xi); y, d\iota_y^T(\xi')) \mid (x, \xi; y, \xi') \in WF(S)\}.$$

Therefore, there must be a point $(x, -\eta; x, \eta) \in WF(S)$ such that $(x, -d\iota_x^T(\eta); x, d\iota_x^T(\eta)) = (x, -d\iota_x^T(\xi); x, d\iota_x^T(\xi))$. Notice, however, that the transpose of $d\iota$ is nothing else than restricting the dual vector ξ to the tangent space of C . Because $WF(S) \subset WF(E)$ there are only two possibilities: $\eta = \xi$ or $\eta = \xi - 2(\xi_a n_0^a)n_0$. The first contradicts our assumption, so we have $\eta = \xi - 2(\xi_a n_0^a)n_0$. Now $(x, -\eta; x, \eta) \in WF(S)$ must hold for every normalised, time-like, future pointing vector $n_0 \in T_x M$. Choosing a sequence of vectors n_0 such that $\eta \rightarrow \xi$ and using the closedness of the wave front set we find again $(x, -\xi; x, \xi) \in WF(S)$. Hence, $WF(E) = WF(S)$. $\square$

B Proof of Theorem 4.18

The computations involved in the proof of Theorem 4.18 are somewhat similar to the computation of the stress-energy-momentum tensor. We will work in components and in local coordinates on O , using Greek indices to indicate the coordinate frame and coordinate derivatives. To ease the notation we will drop the subscript ϵ on the local frame e_a^μ .

As γ^a is independent of ϵ we may use equations (5) to vary

$$\nabla v = \left(\partial_a v - \frac{1}{4} \Gamma_{ab}^c v \gamma_c \gamma^b \right) \gamma^a = e_a^\alpha \left(\partial_\alpha v + \frac{1}{4} e_b^\beta \left\{ \partial_\alpha e_\beta^c - e_\gamma^c \Gamma_{\alpha\beta}^\gamma \right\} v \gamma_c \gamma^b \right) \gamma^a, \quad (22)$$

which yields:

$$\begin{aligned} \delta \nabla v &= \delta e_a^\alpha e_\alpha^d \nabla_d v \gamma^a - \frac{1}{4} \delta e_b^\beta e_\beta^d \Gamma_{ad}^c v \gamma_c \gamma^b \gamma^a + \frac{1}{4} \partial_a \delta e_\beta^c e_b^\beta v \gamma_c \gamma^b \gamma^a \\ &\quad - \frac{1}{4} \delta e_\gamma^c e_a^\alpha e_b^\beta \Gamma_{\alpha\beta}^\gamma v \gamma_c \gamma^b \gamma^a - \frac{1}{4} \delta \Gamma_{\alpha\beta}^\gamma e_a^\alpha e_b^\beta e_\gamma^c v \gamma_c \gamma^b \gamma^a. \end{aligned} \quad (23)$$

We can perform an integration by parts as follows:

$$\begin{aligned} \frac{1}{4} \partial_a \delta e_\beta^c e_b^\beta v \gamma_c \gamma^b \gamma^a &= \frac{-i}{4} P_c (\delta e_\beta^c e_b^\beta v \gamma_c \gamma^b) + \frac{i}{4} \delta e_\beta^c e_b^\beta P_c (v \gamma_c \gamma^b) \\ &\quad - \frac{1}{4} \delta e_\beta^c \partial_a e_b^\beta v \gamma_c \gamma^b \gamma^a - \frac{1}{4} \delta e_\beta^d e_b^\beta \Gamma_{ad}^c v \gamma_c \gamma^b \gamma^a + \frac{1}{4} \delta e_\beta^c e_d^\beta \Gamma_{ab}^d v \gamma_c \gamma^b \gamma^a \\ &= \frac{-i}{4} P_c (\delta e_\beta^c e_b^\beta v \gamma_c \gamma^b) + \frac{i}{4} \delta e_\beta^c e_b^\beta (P_c v) \gamma_c \gamma^b - \frac{1}{4} \delta e_\beta^c e_b^\beta \nabla_a v [\gamma_c \gamma^b, \gamma^a] \\ &\quad - \frac{1}{4} \delta e_\beta^c \partial_a e_b^\beta v \gamma_c \gamma^b \gamma^a + \frac{1}{4} \delta e_b^\beta e_\beta^d \Gamma_{ad}^c v \gamma_c \gamma^b \gamma^a + \frac{1}{4} \delta e_\beta^c e_d^\beta \Gamma_{ab}^d v \gamma_c \gamma^b \gamma^a. \end{aligned} \quad (24)$$

Because $[\gamma_c \gamma^b, \gamma^a] = \gamma_c \{ \gamma^b, \gamma^a \} - \{ \gamma_c, \gamma^a \} \gamma^b = 2\eta^{ab} \gamma_c - 2\delta_c^a \gamma^b$ and $e_\beta^c = g_{\mu\beta} \eta^{cd} e_d^\mu$ we can write:

$$\begin{aligned} -\frac{1}{4} \delta e_\beta^c e_b^\beta \nabla_a v [\gamma_c \gamma^b, \gamma^a] &= -\frac{1}{2} \delta (g_{\mu\beta} \eta^{cd} e_d^\mu) e_b^\beta \eta^{ab} \nabla_a v \gamma_c + \frac{1}{2} \delta e_\beta^c e_b^\beta \nabla_c v \gamma^b \\ &= -\frac{1}{2} \delta g_{\mu\beta} \eta^{cd} e_d^\mu e_b^\beta \eta^{ab} \nabla_a v \gamma_c - \delta e_d^\mu e_\mu^a \nabla_a v \gamma^d \\ &= \frac{1}{2} \delta g^{\alpha\beta} e_\alpha^a e_\beta^b \nabla_a v \gamma_b - \delta e_\alpha^a e_\alpha^d \nabla_d v \gamma^a. \end{aligned} \quad (25)$$

When substituting equations (24) and (25) in (23) we can recombine the terms

$$\frac{-1}{4} \delta e_\beta^c \partial_a e_b^\beta v \gamma_c \gamma^b \gamma^a - \frac{1}{4} \delta e_\gamma^c e_a^\alpha e_b^\beta \Gamma_{\alpha\beta}^\gamma v \gamma_c \gamma^b \gamma^a = \frac{-1}{4} \delta e_\gamma^c e_d^\gamma \Gamma_{ab}^d v \gamma_c \gamma^b \gamma^a$$

to obtain

$$\begin{aligned} \delta \nabla v &= \frac{-i}{4} P_c (\delta e_\beta^c e_b^\beta v \gamma_c \gamma^b) + \frac{i}{4} \delta e_\beta^c e_b^\beta (P_c v) \gamma_c \gamma^b + \frac{1}{2} \delta g^{\alpha\beta} e_\alpha^a e_\beta^b \nabla_a v \gamma_b \\ &\quad - \frac{1}{4} \delta \Gamma_{\alpha\beta}^\gamma e_a^\alpha e_b^\beta e_\gamma^c v \gamma_c \gamma^b \gamma^a. \end{aligned} \quad (26)$$

Note that the variations of the frame δe_a^α cancel out, except in the terms with P_c . These are harmless when we compute $B_0(\delta \nabla S_0 f)$, because both B_0 and v solve the Dirac equation. Therefore, the final answer will not depend on variations of the frame, as desired.

In the last term of equation (26) we can use the symmetry of the Christoffel symbol:

$$\begin{aligned} -\frac{1}{4} \delta \Gamma_{(\alpha\beta)}^\gamma e_a^\alpha e_b^\beta e_\gamma^c v \gamma_c \gamma^b \gamma^a &= -\frac{1}{4} \delta \Gamma_{\alpha\beta}^\gamma e_a^\alpha e_b^\beta e_\gamma^c v \gamma_c \eta^{ab} = -\frac{1}{4} \delta \Gamma_{\alpha\beta}^\gamma g^{\alpha\beta} e_\gamma^c v \gamma_c \\ &= -\frac{1}{4} \delta g^{\gamma\mu} g_{\mu\nu} \Gamma_{\alpha\beta}^\nu e_\gamma^c v \gamma_c - \frac{1}{4} \partial_\alpha \delta g_{\beta\mu} e_a^\mu g^{\alpha\beta} v \gamma^a \\ &\quad + \frac{1}{8} \partial_\mu \delta g_{\alpha\beta} e_a^\mu g^{\alpha\beta} v \gamma^a \end{aligned} \quad (27)$$

We handle the last term using an integration by parts as before:

$$\begin{aligned} \frac{1}{8} \partial_a \delta g_{\alpha\beta} g^{\alpha\beta} v \gamma^a &= \frac{-i}{8} P_c (\delta g_{\alpha\beta} g^{\alpha\beta} v) + \frac{i}{8} \delta g_{\alpha\beta} g^{\alpha\beta} P_c v - \frac{1}{8} \delta g_{\alpha\beta} \partial_a g^{\alpha\beta} v \gamma^a \\ &= \frac{-i}{8} P_c (\delta g_{\alpha\beta} g^{\alpha\beta} v) + \frac{i}{8} \delta g_{\alpha\beta} g^{\alpha\beta} P_c v - \frac{1}{8} \delta g^{\alpha\beta} \partial_a g_{\alpha\beta} v \gamma^a, \end{aligned} \quad (28)$$

where we used $\delta g_{\alpha\beta} \partial_a g^{\alpha\beta} = -\delta g^{\alpha\beta} g_{\alpha\mu} g_{\beta\nu} \partial_a g^{\mu\nu} = \delta g^{\alpha\beta} \partial_a g_{\alpha\beta}$. The penultimate term in (27) is:

$$\begin{aligned}
-\frac{1}{4} \partial_\alpha \delta g_{\beta\mu} e_a^\mu g^{\alpha\beta} v \gamma^a &= \frac{1}{4} \partial_b (\delta g^{\alpha\beta} g_{\alpha\mu} g_{\beta\nu}) e_a^\mu e_\rho^b g^{\rho\nu} v \gamma^a \\
&= \frac{1}{4} \partial_b (\delta g^{\alpha\beta} e_\alpha^a e_\beta^b) v \gamma_a - \frac{1}{4} \delta g^{\alpha\beta} g_{\alpha\mu} g_{\beta\nu} \partial_b (e_a^\mu e_\rho^b g^{\rho\nu}) v \gamma^a \\
&= \frac{1}{4} \nabla_b (\delta g^{\alpha\beta} e_\alpha^a e_\beta^b) v \gamma_a - \frac{1}{4} \delta g^{\alpha\beta} (\Gamma_{bc}^a e_\alpha^c e_\beta^b + \Gamma_{bc}^b e_\alpha^a e_\beta^c) v \gamma_a \\
&\quad - \frac{1}{4} \delta g^{\alpha\beta} g_{\alpha\mu} g_{\beta\nu} \partial_b (e_a^\mu e_\rho^b g^{\rho\nu}) v \gamma^a.
\end{aligned} \tag{29}$$

The first term on the right-hand side of equation (29) is

$$\frac{1}{4} \nabla_b (\delta g^{\alpha\beta} e_\alpha^a e_\beta^b) v \gamma_a = \frac{1}{4} \nabla_b (\delta g^{\alpha\beta} e_\alpha^a e_\beta^b v \gamma_a) - \frac{1}{4} \delta g^{\alpha\beta} e_\alpha^a e_\beta^b \nabla_b v \gamma_a. \tag{30}$$

The other terms can be simplified with some computation:

$$\begin{aligned}
&-\frac{1}{4} \delta g^{\alpha\beta} (\Gamma_{bc}^a e_\alpha^c e_\beta^b + \Gamma_{bc}^b e_\alpha^a e_\beta^c + g_{\alpha\mu} g_{\beta\nu} \eta^{ac} \partial_b (e_c^\mu e_\rho^b g^{\rho\nu})) v \gamma_a \\
&= -\frac{1}{4} \delta g^{\alpha\beta} \left(-\partial_\beta e_\alpha^a + e_\gamma^\alpha \Gamma_{\beta\alpha}^\gamma - e_\alpha^a \partial_c e_\beta^c + e_\alpha^a \Gamma_{\mu\beta}^\mu + e_\alpha^a g_{\beta\nu} \partial_\rho g^{\rho\nu} + e_\alpha^a \partial_b e_\beta^b + g_{\alpha\mu} \eta^{ac} \partial_\beta e_c^\mu \right) v \gamma_a \\
&= -\frac{1}{4} \delta g^{\alpha\beta} \left(-\eta^{ac} e_c^\mu \partial_\beta g_{\alpha\mu} + e_\gamma^\alpha \Gamma_{\beta\alpha}^\gamma + e_\alpha^a \Gamma_{\mu\beta}^\mu - e_\alpha^a g^{\rho\nu} \partial_\rho g_{\beta\nu} \right) v \gamma_a \\
&= -\frac{1}{8} \delta g^{\alpha\beta} (-2e_\gamma^\alpha g^{\gamma\mu} \partial_\beta g_{\alpha\mu} + e_\gamma^\alpha g^{\gamma\mu} (2\partial_\beta g_{\alpha\mu} - \partial_\mu g_{\alpha\beta}) + e_\alpha^a g^{\mu\gamma} \partial_\beta g_{\mu\gamma} - 2e_\alpha^a g^{\rho\nu} \partial_\rho g_{\beta\nu}) v \gamma_a \\
&= \frac{1}{8} \delta g^{\alpha\beta} (e_\gamma^\alpha g^{\gamma\mu} \partial_\mu g_{\alpha\beta} + 2e_\alpha^a g_{\beta\mu} g^{\rho\nu} \Gamma_{\rho\nu}^\mu) v \gamma_a.
\end{aligned} \tag{31}$$

Substituting equations (27-31) into (26) yields:

$$\begin{aligned}
\delta \nabla v &= \frac{-i}{4} P_c (\delta e_\beta^c e_b^\beta v \gamma_c \gamma^b) + \frac{i}{4} \delta e_\beta^c e_b^\beta (P_c v) \gamma_c \gamma^b - \frac{i}{8} P_c (\delta g_{\alpha\beta} g^{\alpha\beta} v) + \frac{i}{8} \delta g_{\alpha\beta} g^{\alpha\beta} P_c v \\
&\quad + \frac{1}{4} \delta g^{\alpha\beta} e_\alpha^a e_\beta^b \nabla_a v \gamma_b + \frac{1}{4} \nabla_b (\delta g^{\alpha\beta} e_\alpha^a e_\beta^b v \gamma_a).
\end{aligned} \tag{32}$$

Using Lemma 4.17 we find for a spinor $u \in C^\infty(DM)$:

$$\begin{aligned}
\delta \nabla u &= \frac{i}{4} P (\delta e_\beta^c e_b^\beta \gamma^b \gamma_c u) - \frac{i}{4} \delta e_\beta^c e_b^\beta \gamma^b \gamma_c (P u) + \frac{i}{8} P (\delta g_{\alpha\beta} g^{\alpha\beta} u) - \frac{i}{8} \delta g_{\alpha\beta} g^{\alpha\beta} P u \\
&\quad + \frac{1}{4} \delta g^{\alpha\beta} e_\alpha^a e_\beta^b \gamma_b \nabla_a u + \frac{1}{4} \nabla_b (\delta g^{\alpha\beta} e_\alpha^a e_\beta^b \gamma_a u).
\end{aligned} \tag{33}$$

Using Proposition 4.16 and equations (32,33) we notice that the terms with P_c and P cancel out in the following equality, because B_0 and $S_0 f$ both satisfy the Dirac equation:

$$\begin{aligned}
\delta(\beta_\epsilon B_0(f)) &= -B_0(\delta P_\epsilon S_0 f) = \frac{i}{4} B_0(\delta g^{\alpha\beta} e_\alpha^a e_\beta^b \gamma_b \nabla_a S_0 R f) + \frac{i}{4} B_0(\nabla_b (\delta g^{\alpha\beta} e_\alpha^a e_\beta^b \gamma_a S_0 R f)) \\
&= \frac{i}{4} \delta g^{\alpha\beta} e_\alpha^a e_\beta^b (B_0(\gamma_b \nabla_a S_0 R f) - \nabla_b (B_0(\gamma_a S_0 R f))).
\end{aligned} \tag{34}$$

We now compare with Proposition 4.14 to get the final result.

References

- [1] H. Araki, *On the diagonalization of a bilinear Hamiltonian by a Bogoliubov transformation*, Publ. Res. Inst. Math. Sci. Ser. A **4** (1968/1969) 387–412

- [2] C. Bär, N. Ginoux and F. Pfäffle, *Wave equations on Lorentzian manifolds and quantization*, EMS, Zürich, (2007)
- [3] O. Bratteli and D.W. Robinson, *Operator algebras and quantum statistical mechanics 2*, Springer, Berlin (1996)
- [4] R. Brunetti, K. Fredenhagen and R. Verch, *The generally covariant locality principle—a new paradigm for local quantum field theory*, Commun. Math. Phys. **237** (2003), 31–68
- [5] R. Brunetti, K. Fredenhagen and M. Köhler, *The microlocal spectrum condition and Wick polynomials of free fields on curved spacetimes*, Commun. Math. Phys. **180** (1996), 633–652
- [6] D. Canarutto and A. Jadczyk, *Fundamental geometric structures for the Dirac equation in general relativity*, Acta Appl. Math. **51** (1998), 59–92
- [7] Y. Choquet-Bruhat, C. de Witt-Morette and M. Dillard-Bleick, *Analysis, manifolds and physics*, North Holland, Amsterdam (1977)
- [8] R. Coquereaux, *Clifford algebras, spinors and fundamental interactions: Twenty Years After*, arXiv:math-ph/0509040v1 (2005)
- [9] C. D’Antoni and S. Hollands, *Nuclearity, local quasiequivalence and split property for Dirac quantum fields in curved spacetime*, Commun. Math. Phys. **261** (2006), 133–159
- [10] C. Dappiaggi, T.-P. Hack and N. Pinamonti, *The extended algebra of observables for Dirac fields and the trace anomaly of their stress-energy tensor*, arXiv:0904.0612 [math-ph]
- [11] S.P. Dawson and C.J. Fewster, *An explicit quantum weak energy inequality for Dirac fields in curved spacetimes*, Class. Quantum Grav. **23** (2006), 6659–6681
- [12] N. Dencker, *On the propagation of polarization sets for systems of real principal type*, J. Funct. Anal. **46** (1982), 351–372
- [13] J. Dieudonné, *Treatise on analysis, Vol. III*, Academic Press, New York-London (1972)
- [14] J. Dimock, *Dirac quantum fields on a manifold*, Trans. Amer. Math. Soc. **269** (1982), 133–147
- [15] J.J. Duistermaat and L. Hörmander, *Fourier integral operators. II*, Acta Math. **128** (1972), 183–269
- [16] C.J. Fewster, *Quantum energy inequalities and local covariance II: categorical formulation*, Gen. Relativ. Gravit. **39** (2007) 18551890
- [17] C.J. Fewster and R. Verch, *A quantum weak energy inequality for Dirac fields in curved spacetime*, Commun. Math. Phys. **225** (2002), 331–359
- [18] M. Forger and H. Römer, *Currents and the energy-momentum tensor in classical field theory: a fresh look at an old problem*, Ann. Phys. **309** (2004), 306–389
- [19] R. Geroch, *Spinor structures of space-times in general relativity. I*, J. Mathematical Phys. **9** (1968), 1739–1744
- [20] R. Geroch, *Spinor structures of space-times in general relativity. II*, J. Mathematical Phys. **11** (1970), 343–348
- [21] R.H. Good Jr., *Properties of the Dirac matrices* Rev. Mod. Phys. **27** (1955) 187–211
- [22] S. Hollands, *The Hadamard condition for Dirac fields and adiabatic states on Robertson-Walker spacetimes*, Commun. Math. Phys. **216** (2001) 635–661
- [23] S. Hollands, *The operator product expansion for perturbative quantum field theory in curved spacetime*, Commun. Math. Phys. **273** (2007) 1–36

- [24] L. Hörmander, *The Analysis of Linear Partial Differential Operators I*, Springer, Berlin (2003)
- [25] S. Kobayashi and K. Nomizu, *Foundations of differential geometry* vol I, Interscience, New York, (1963)
- [26] K. Kratzert, *Singularity structure of the two point function of the free Dirac field on a globally hyperbolic spacetime*, Ann. Phys. (8) **9** (2000), 475–498
- [27] H.B. Lawson and M.-L. Michelson, *Spin geometry*, Princeton university press, Princeton, 1989
- [28] A. Lichnerowicz, *Champs spinoriels et propagateurs en relativité générale*, Bull. Soc. Math. France **92** (1964) 11–100
- [29] S. Mac Lane, *Categories for the working mathematician*, Springer, New York, (1971)
- [30] S. Mac Lane and I. Moerdijk, *Sheaves in geometry and logic: a first introduction to topos theory*, Springer, New York, (1992)
- [31] W. Pauli, *Contributions mathématiques à la théorie des matrices de Dirac*, Ann. Inst. H. Poincaré **6** (1936), 109–136
- [32] M.J. Radzikowski, *Micro-Local Approach to the Hadamard Condition in Quantum Field Theory on Curved Space-Time*, Commun. Math. Phys. **179** (1996), 529–553
- [33] J. Roberts, G. Ruzzi, *A cohomological description of connections and curvature tensors over posets*, Theory and Applications of Categories **16** (2006), 855–895
- [34] H. Sahlmann and R. Verch, *Microlocal spectrum condition and Hadamard form for vector-valued quantum fields in curved spacetime*, Rev. Math. Phys. **13** (2001), 1203–1246
- [35] K. Sanders, *Equivalence of the (generalised) Hadamard and microlocal spectrum condition for (generalised) free fields in curved spacetime*, Commun. Math. Phys. (DOI) 10.1007/s00220-009-0900-7
- [36] K. Sanders, *Aspects of locally covariant quantum field theory*, PhD thesis, University of York, also available online, arXiv: 0809.4828v1[math-ph]
- [37] K. Schmüdgen, *Unbounded operator algebras and representation theory*, Birkhäuser Verlag, Basel (1990)
- [38] G. Segal, *Classifying spaces and spectral sequences.*, Inst. Hautes Études Sci. Publ. Math. **34** (1968), 105–112
- [39] A. Strohmaier, R. Verch and M. Wollenberg, *Microlocal analysis of quantum fields on curved space-times: analytic wave front sets and Reeh-Schlieder theorems*, J. Math. Phys. **43** (2002), 5514–5530
- [40] J. Tolksdorf, *Clifford modules and generalized Dirac operators*, Internat. J. Theoret. Phys. **40** (2001), 191–209
- [41] B.L. van der Waerden, *Group theory and quantum mechanics*, Springer, Berlin (1974)
- [42] R.M. Wald, *General relativity*, The University of Chicago Press, Chicago and London (1984)